

## Allelopathic activity of decomposing straw of wheat and oat and associated soil on some crop species

L.S. Dias

*Departamento de Biologia, Universidade de Évora, Ap 94, P-7001 Évora Codex, Portugal*

(Accepted for publication 9 August 1990)

### ABSTRACT

Dias, L.S., 1991. Allelopathic activity of decomposing straw of wheat and oat and associated soil on some crop species. *Soil Tillage Res.*, 21: 113–120.

Allelopathic effects of decomposing straw of wheat (*Triticum aestivum* L.) and oat (*Avena sativa* L.) and associated soil were assessed on germination and early root growth of wheat, oat, and subterranean clover (*Trifolium brachycalycinum* Katzn. & Morley). Although allelopathic activity was widespread during decomposition, only speed of germination and root length were affected with both inhibitory and stimulatory effects. Important differences in sensitivity were found among the target species, with mainly subterranean clover being stimulated, and the grasses mainly inhibited. Conversely, only minor differences in effectiveness were found between the source species. Soil treatments were less effective than straw treatments, with a generalised pattern of inhibitory cycles being evident. Results suggest that allelopathic activity might be of minor relevance in mulch tillage involving the investigated species. However, further research to characterise and identify the chemicals responsible for observed allelopathic effects is needed.

Pedidos de cópia desta publicação para Luís Silva Dias, Departamento de Biologia, Universidade de Évora, Ap. 94, 7002-554 Évora, Portugal ou, de preferência, para [lsdias@uevora.pt](mailto:lsdias@uevora.pt).

Reprint requests to Luís Silva Dias, Departamento de Biologia, Universidade de Évora, Ap. 94, 7002-554 Évora, Portugal or preferably to [lsdias@uevora.pt](mailto:lsdias@uevora.pt).