



## Relative timing of transcurrent displacements in northern Gondwana: U–Pb laser ablation ICP–MS zircon and monazite geochronology of gneisses and sheared granites from the western Iberian Massif (Portugal)

M.F. Pereira <sup>a,\*</sup>, J.B. Silva <sup>b</sup>, K. Drost <sup>c</sup>, M. Chichorro <sup>d</sup>, A. Apraiz <sup>e</sup>

<sup>a</sup> Departamento de Geociências, CGE, Escola de Ciências e Tecnologia, Universidade de Évora, Portugal

<sup>b</sup> Departamento de Geologia, Faculdade de Ciências, IDL, Universidade de Lisboa, Edifício C3, Campo Grande, Lisboa, Portugal

<sup>c</sup> Department of Earth Sciences, University of Cape Town, AEON Earth Lab, CGE, South Africa

<sup>d</sup> CICEGe, Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, Quinta da Torre, 2829-516 Caparica, Portugal

<sup>e</sup> Geodinamika Saila, Zientzia eta Teknologia Fak., Euskal Herriko Unibertsitatea. Apt. 644, 48080 Bilbo, Spain

### ARTICLE INFO

#### Article history:

Received 1 May 2009

Received in revised form 8 August 2009

Accepted 26 August 2009

Available online 10 September 2009

#### Keywords:

Transcurrent deformation  
Variscan orogeny  
Pangaea assembly  
Northern Gondwana  
Coimbra–Córdoba shear zone  
Porto–Tomar fault zone

### ABSTRACT

The Variscan belt of Western and Central Europe was formed by the oblique subduction of the Rheic Ocean and the collision of Laurussia with Gondwana during the Late Palaeozoic. We present field relationships and new U–Pb LA-ICP-MS zircon and monazite ages for Variscan gneisses and granites from a key section of the western Iberian Massif. The Martinchel section records the interplay of two kilometre-scale Variscan transcurrent shear zones active in the Gondwana basement of Pangaea: the Porto–Tomar fault zone (PTFZ) and the Coimbra–Córdoba shear zone (CCSZ). Different kinematic models have been invoked to explain the formation and evolution of these major Variscan structures mainly based on assumptions made in the absence of reliable radiometric ages. We show that: (1) ductile deformation and metamorphism were active in the CCSZ during the Visean–Serpukhovian (c.335–318 Ma) and created conditions for amphibolite facies metamorphism and coeval emplacement of granites; and (2) later ductile–brittle deformation related to dextral movements along the PTFZ overprinted the earlier foliation and folds derived from the CCSZ deformation, and deformed the previously intruded granites. U–Pb dating of zircon and monazites yield c.335 Ma ages for the ductile deformation developed under amphibolite facies metamorphic conditions in the Martinchel gneisses of the CCSZ. The gneisses were intruded by granites at c.335–318 Ma, and both were later deformed under ductile–brittle conditions by dextral motion on the PTFZ. The geometry of the Martinchel gneisses (typical of the CCSZ) changed from one of thrusting to one of normal faulting by refolding of the early foliation, stretching lineation and asymmetric structures related to the later PTFZ dextral shear episode. This pattern of interference is not fully considered in previous models and may lead to incorrect tectonic interpretations. According to our data and recently published ages, we suggest that the PTFZ was active after the Serpukhovian–Kasimovian since the c.318–308 Ma granites are deformed by north–south (170°) dextral shear planes. These data are critical to the interpretation of large-scale Carboniferous transcurrent displacements in northern Gondwana (Iberian Massif), and bear upon global models of crustal deformation that emphasize the importance of long-lived dextral movements during the collision between northern Gondwana and Laurussia following the closure of the Rheic Ocean.

© 2009 International Association for Gondwana Research. Published by Elsevier B.V. All rights reserved.

### 1. Introduction

The Variscan belt of Western and Central Europe is a part of a large-scale orogenic system, with continuations into North America (Ouachitas and southern Appalachians) and West Africa (Mauritanides) (Matte, 1991; Shelley and Bossiere, 2000; Simancas et al., 2004;

Martinez-Catalan et al., 2007; Melleton et al., 2010; Murphy et al., 2009). In the 1970s and early 1980s, a number of structural schemes were proposed to clarify the complex architecture of the Variscan belt of Western Europe with respect to the subduction of the Rheic Ocean and the collision of Laurentia–Baltica with Gondwana (Arthaud and Matte, 1977; Bard et al., 1980; Matte, 1986). Based on analogies with Alpine and Himalayan belts, these authors attempted to explain the time and space constraints of deformation, metamorphism and magmatism linked to the evolution of this Palaeozoic belt.

Some of the most prominent structures in the Variscan belt of Western and Central Europe are major fracture zones in the Precambrian

\* Corresponding author.

E-mail addresses: [mpereira@uevora.pt](mailto:mpereira@uevora.pt) (M.F. Pereira), [jbsilva@fc.ul.pt](mailto:jbsilva@fc.ul.pt) (J.B. Silva), [kerstin.drost@geo.uib.no](mailto:kerstin.drost@geo.uib.no) (K. Drost), [ma.chichorro@fct.unl.pt](mailto:ma.chichorro@fct.unl.pt) (M. Chichorro), [arturo.apraiz@ehu.es](mailto:arturo.apraiz@ehu.es) (A. Apraiz).