

A Greenhouse Tomato Crop Grey Mould Disease Early Warning System

M. Neto¹, F. Baptista², L. M. Navas³ and G. Ruiz³

¹ISEGI-UNL, Campus de Campolide, 1070-312 Lisbon, Portugal

²Department of Agricultural Engineering, University of Évora/ECT/ICAAM, ZIP 94, 7002-554 Évora, Portugal

³Department of Agricultural and Forestry Engineering, University of Valladolid. Avda. Madrid, 44, 34004 Palencia, Spain
mneto@isegi.unl.pt

Abstract

Tomato is a very important crop in the Mediterranean region in general and in Portugal in particular being the production for fresh consumption made essentially in greenhouses. *Botrytis cinerea* Pers.: Fr. is the causal agent of grey mould disease and is one of the most important diseases affecting greenhouse tomato crops, high relative humidity and the presence of free water on the plant surfaces have been recognized as favourable to the development of this disease.

The availability of a early warning system providing to the tomato grower alerts with information of the potential favoured conditions for the disease appearance in its early stages or even before can have a very positive impact in reducing the economic and environmental impacts due to a more rational and efficient disease control management.

Today we have the necessary technology to build and launch an Internet based early warning system for grey mould disease in greenhouse tomato crop supported by a wireless sensor network. In this paper a prototype for such a system is presented.

From the research conducted until the moment the proposed solution is viable and the next step will be to validate it in the field in different locations and with distinct greenhouses conditions.

Keywords: Greenhouse tomato; grey mould; wireless sensor network; early warning system

Introduction

In the Mediterranean region greenhouse areas are significant with tomato being the most commonly grown vegetable (Castilla, 2002). In Portugal according to available statistical information greenhouse tomato crop was cultivated in 1,550 ha with a total production of 98,646 ton in 2003 (GPP, 2007).

Botrytis cinerea Pers.: Fr. is the causal agent of grey mould disease and is one of the most important diseases affecting greenhouse tomato crops. This disease could be responsible for production losses of 20% and treatments could represent about 60% of the total fungicides used over a cropping season (Prieto *et al.*, 2003). Conditions inside greenhouses; warm, humid, high plant density and frequent handling are conducive to the establishment and spread of the pathogen.