

## XVII Congresso Nacional de Bioquímica

Poster Abstract Submission

Name: Ana Rodrigues Costa.....

Research Group: ICAAM-CTA (Grupo de Ciência e Tecnologia Animal do ICAAM-Univ. Évora) ...

E-mail: acrc@uevora.pt.....

### Abstract

**Title:** Implication of AMPK in glucose-evoked modulation of Na,K-ATPase

**Authors:** Ana R. Costa<sup>1,2</sup>, Célia M Antunes<sup>1,2,3</sup>, Júlio Cruz-Morais<sup>1,2</sup>

**Affiliation:** 1-ICAAM; 2 – Departamento de Química, Univ. Évora; 3 – CNC-Coimbra

**Themes:** Clinical Biochemistry and Mechanisms of Disease.....

**Type of Presentation:** Poster

**Text:**

**Background and aims:** Na,K-ATPase is an integral membrane protein that maintains the gradients of Na<sup>+</sup> and K<sup>+</sup>, using the energy of ATP hydrolysis, maintaining the ionic gradients that allow electrical activity to occur. It has been demonstrated that, in pancreatic  $\beta$ -cells, Na,K-ATPase is regulated by glucose and that this phenomenon is impaired in glucose intolerant subjects. However, the mechanism underlying glucose-induced modulation of Na,K-ATPase is still unclear.

The AMP-activated protein kinase (AMPK) is a molecular key player in energy homeostasis, providing exquisite sensitivity to small changes in intracellular AMP levels and thus to intracellular [ATP]/[ADP] ratio, that is known to activate protein regulatory pathways. Since in pancreatic  $\beta$ -cell, glucose has marked effects on oxidative metabolism and total intracellular ATP and AMP levels, the involvement of AMPK in the cascade of events regulating Na,K-ATPase regulation in pancreatic  $\beta$ -cells was postulated. The aim of this work was to evaluate the putative role of AMPK in the glucose-evoked regulation of Na,K-ATPase activity in the pancreatic  $\beta$ -cell.

**Materials and methods:** Pancreatic  $\beta$ -cells from normal (control) or glucose-intolerant Wistar rats (GIR) were isolated and cultured (48h). Cell batches were pre-incubated (30min) with 2.1mM glucose to reach basal activity. Afterwards cells were challenged to 8.4mM glucose for 20min, in the presence or absence of AMPK agonists (AICAR) and antagonists (compound C; CC). ATPase activity was assessed in intact cells by colorimetric quantification of Pi formed in 30min. Na,K-ATPase activity was calculated by the difference between the activities obtained in the absence and in presence the of 1mM ouabain.

**Results:** In basal conditions the activity of Na,K-ATPase from normal and GIR pancreatic  $\beta$ -cell was similar ( $0.184 \pm 0.030$  and  $0.186 \pm 0.020$   $\mu\text{molPi}/\text{min}/\text{mgProt}$ , respectively). Challenging the control  $\beta$ -cells with glucose 8.4mM evoked a 62% reduction of Na,K-ATPase activity whereas in GIR  $\beta$ -cells a significantly lower inhibition (40%) was observed. The addition of AICAR 1mM abolished glucose-induced Na,K-ATPase inhibition ( $0.166 \pm 0.011$   $\mu\text{molPi}/\text{min}/\text{mg}$ ). In control  $\beta$ -cell, the addition of CC 10  $\mu\text{M}$  had no effect on glucose-induced inhibition of Na,K-ATPase. In the contrary, in GIR  $\beta$ -cells it significantly potentiated glucose-evoked inhibition of Na,K-ATPase reaching values similar to that

# XVII CONGRESSO NACIONAL DE BIOQUÍMICA

15, 16 e 17 DEZEMBRO 2010, PORTO

observed in the controls (66%).

**Conclusions:** The AMPK agonist AICAR counteracts the inhibitory action of glucose on Na,K-ATPase of control  $\beta$ -cells whereas CC amplified the glucose-induced inhibition of Na,K-ATPase in GIR  $\beta$ -cells. These results suggest that AMPK plays a central role in the cascade of events underlying glucose-induced modulation of Na,K-ATPase and that the defect must be upstream of AMPK. Finally, abnormal glucose-induced regulation of Na,K-ATPase occurs prior to overt type 2 diabetes and might be a feature in the disease development.