

Comparative Biochemistry and Physiology, Part A 189 (2015) 67–75

Structural lipid changes and Na⁺/K⁺-ATPase activity of gill cells' basolateral membranes during saltwater acclimation in sea lamprey (*Petromyzon marinus*, L.) juveniles

Maria João Lança^a, Maria Machado^b, Ana Filipa Ferreira^c, Bernardo Ruivo Quintella^{c,d}, Pedro Raposo de Almeida^c

^a Departamento de Zootecnia, Escola de Ciências e Tecnologia, ICAAM–Instituto de Ciências Agrárias e Ambientais Mediterrânicas, Universidade de Évora, Largo dos Colegiais 2, Évora 7004-516, Portugal

^b Hospital Veterinário das Laranjeiras, Rua São Tomás de Aquino 8C, Lisboa 100-203, Portugal

^c MARE–Centro de Ciências do Mar e do Ambiente, Departamento de Biologia, Escola de Ciências e Tecnologia, Universidade de Évora, Largo dos Colegiais 2, Évora 7004-516, Portugal

^d Departamento de Biologia Animal, Faculdade de Ciências da Universidade de Lisboa, Campo Grande, Lisboa 1749-016, Portugal

article info abstract

Article history:

Received 3 June 2015

Received in revised form 29 July 2015

Accepted 29 July 2015

Available online 2 August 2015

Keywords: basolateral membrane; gill Cell; juveniles; *Petromyzon marinus*; phospholipids' composition; saltwater acclimation

Seawater acclimation is a critical period for anadromous species and a process yet to be understood in lampreys. Considering that changes in lipid composition of the gill cells' basolateral membranes may disrupt the major transporter Na⁺K⁺-ATPase, the goal of this study was to detect changes at this level during juvenile sea lamprey seawater acclimation. The results showed that saltwater acclimation has a direct effect on the fatty acid composition of gill cells basolateral membrane's phospholipids. When held in full-strength seawater, the fatty acid profile of basolateral membrane's phospholipids suffered a restructure by increasing either saturation or the ratio between oleic acid and eicosapentaenoic acid. Simultaneously, the activity of Na⁺K⁺-

ATPase revealed a significant and positive correlation with basolateral membrane's cholesterol content in the presence of highest salinity. Our results pointed out for lipid adjustments involving the functional transporter present on the gill cell basolateral membranes to ensure the role played by branchial Na^+K^+ -ATPase in ion transport during saltwater acclimation process. The responses observed contributed to the strategy adopted by gill cell's basolateral membranes to compensate for osmotic and ionic stressors, to ensure the success of the process of seawater acclimation associated with the downstream trophic migration of juvenile sea lamprey.

© 2015 Elsevier Inc. All rights reserved.