

## Diet and disease in Tomar, Portugal: Comparing stable carbon and nitrogen isotope ratios between skeletons with and without signs of infectious disease



Ana Curto<sup>a,b,\*</sup>, Patrick Mahoney<sup>a</sup>, Anne-France Maurer<sup>c</sup>, Cristina Barrocas-Dias<sup>c,d</sup>,  
Teresa Fernandes<sup>b,e</sup>, Geraldine E. Fahy<sup>a</sup>

<sup>a</sup> Human Osteology Lab, Skeletal Biology Research Centre, School of Anthropology and Conservation, University of Kent, Canterbury, UK

<sup>b</sup> Research Centre for Anthropology and Health, Department of Life Sciences, University of Coimbra, Coimbra, Portugal

<sup>c</sup> HERCULES Laboratory, University of Évora, Largo Marquês de Mariaiva 8, 7000-809, Évora, Portugal

<sup>d</sup> School of Sciences and Technology, Chemistry Department, University of Évora, Rua Romão Ramalho 59, Évora, Portugal

<sup>e</sup> School of Sciences and Technology, Biology Department, University of Évora, Pólo da Mitra, Apartado 94, Évora, Portugal

### ARTICLE INFO

#### Keywords:

Paleodiet  
Paleopathology  
Periostitis  
Infectious disease

### ABSTRACT

**Objectives:** This study explored the correspondence between stable isotope ratios and indicators of non-specific (periostitis and/or osteomyelitis) and specific (venereal syphilis) disease in a sample of human skeletons from a Portuguese archaeological collection. Additionally, this study examined stable carbon ( $\delta^{13}\text{C}$ ) and nitrogen ( $\delta^{15}\text{N}$ ) isotope ratios between individuals at different disease stages.

**Materials and methods:**  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  data from previously analysed skeletons without signs of infectious disease or physiological stress ( $n = 32$ ) were compared to new data from skeletons with active ( $n = 6$ ), healed ( $n = 7$ ) or a combination of both lesions ( $n = 10$ ). Skeletons with lesions ( $n = 23$ ) were also grouped as having only healed tibial periostitis ( $n = 7$ ), generalised non-specific ( $n = 5$ ) and generalised specific infections ( $n = 2$ ). The skeletons with lesions that did not fit into these groups ( $n = 9$ ) were not used in this analysis.

**Results:** The  $\delta^{15}\text{N}$  from skeletons with non-specific generalised infections in several bones differed significantly when compared to skeletons that had either only healed tibial periostitis or were without lesions. Skeletons with venereal syphilis had similar mean  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  to either skeletons without signs of disease or those with only healed tibial periostitis.

**Discussion:** These results suggest different diets may be linked into an individual's susceptibility to these pathogens. Diet influences resistance to infectious disease, while infections decrease nutrient availability, increase malabsorption and resting energy expenditure. Potentially therefore, combining isotopic evidence of diet with pathology may contribute to a new understanding of health and lifestyle in the past.

## 1. Introduction

### 1.1. Effect of diet on health

Nutritional stress may result in either greater susceptibility to physiological stress or greater resilience to stress later in life (Bogin et al., 2007). Malnutrition impairs the immune system (e.g. Calder, 2013; Calder and Jackson, 2000; Scrimshaw and SanGiovanni, 1997). Individuals with poorer nutrition are less resistant to infectious diseases, and infectious disease decreases nutrient availability (e.g. Martorell, 1980; Mata et al., 1971). The effect of protein-energy malnutrition on aspects of immune function and susceptibility to infection (e.g. Calder and Jackson, 2000; Kuvibidila et al., 1993; Scrimshaw and

SanGiovanni, 1997; Woodward, 1998; Woodward, 2001) affects practically all forms of immunity, in particular cell mediated immunity (Kuvibidila et al., 1993; Woodward, 1998, 2001), immune barrier function (Deitch et al., 1990; Sherman et al., 1985) and the functioning of lymphoid organs (Lee and Woodward, 1996; Woodward and Miller, 1991). On the other hand, infections can decrease nutrient availability due to malabsorption (e.g. Mitra et al., 1997) and increase resting energy expenditure, altering the metabolism and redistribution of nutrients (Calder, 2013). However, if nutrition is adequate, diseases like tuberculosis may have a less severe infection, instead of an exacerbated infection, resulting in prolonged chronic infections with a higher probability to affect the skeleton (Ulijaszek et al., 2012).

\* Corresponding author. Human Osteology Lab, Skeletal Biology Research Centre, School of Anthropology and Conservation, University of Kent, Canterbury, UK.  
E-mail address: [arqc3@kent.ac.uk](mailto:arqc3@kent.ac.uk) (A. Curto).

<https://doi.org/10.1016/j.jas.2019.03.005>

Received 18 July 2018; Received in revised form 12 March 2019; Accepted 12 March 2019

Available online 01 April 2019

0305-4403/© 2019 Published by Elsevier Ltd.

## 1.2. Skeletal lesions as health indicators

Health is a complex state that can be reflected through skeletal indicators of physiological stress (Temple and Goodman, 2014). Physiological stress can be related to a wide variety of factors such as disease and nutritional deficiencies (Armstrong, 2003; Goodman and Martin, 2002; Huss-Ashmore et al., 1992;). Even though systemic physiological stress is not directly observable in the skeleton their consequences, in some cases, are (Klaus, 2014).

Infectious diseases were a significant cause of death in past populations, particularly prior to the antibiotic era (Ortner and Putschard, 1985). Pathogens can reach the skeleton by direct infection through wounds, extensions from adjacent soft tissue infections or spread by the blood from the site of a remote infection (Ortner and Putschard, 1985; Ortner, 2003). The body reacts to infection through an inflammatory response which aims to neutralize the pathogen and repair the resultant damage (Weston, 2012). Infection damages the normal cells and accelerates the cell turnover (inflammatory process) (Ragsdale and Lehmer, 2012). Inflammation affects the bone tissue at some level through the production of pathological skeletal phenotypes (e.g. Ragsdale and Lehmer, 2012; Redlich and Smolen, 2012). However, inflammation can be caused by other factors (e.g. Ortner, 2003; Ortner and Putschard, 1985). Bone reacts in a limited number of ways (production or destruction of bone, or a combination of production and destruction of bone) for either infection or other causes such as trauma (e.g. Ragsdale and Lehmer, 2012; Weston, 2008, 2009). However, by analysing the skeleton as a whole and taking into account other bone-forming disorders, systemic non-specific infection remains a contextually plausible diagnostic option (Klaus, 2014).

The bone changes associated with periostitis, an inflammation of the periosteum resulting in deposition of new bone (Bush, 1989), vary from one or more layers of woven or compact bone to spiculae perpendicular to the surface of the bone (Ortner, 2003). Periostitis not associated with a specific skeletal syndrome, particularly on the tibiae, can be linked to pathogens such as *Staphylococcus* or *Streptococcus* (Goodman and Martin, 2002). However, the periosteum responds in a similar way regardless of the etiology (Weston, 2008, 2009). Tibial periostitis is the most commonly reported skeletal lesions in archaeological samples (e.g. DeWitte, 2010; Weston, 2012), being frequently considered an indicator of non-specific physiological stress (e.g. DeWitte, 2010; Robb et al., 2001).

In case of infection leading to pathological new bone formation, inflammation-derived pathological periosteal new bone formation is rooted in biological stress (Klaus, 2014). Osteomyelitis is the result of the introduction of infectious agents into bone, affecting the medullary cavity (Ortner and Putschard, 1985; Ortner, 2003). Bones with osteomyelitis can present a combination of cloacae, sequestered bone and involucrum or only reactive bone formation in the marrow and outer cortex that can result in smooth or lumpy compact bone (Ortner and Putschard, 1985; Ortner, 2003; Pinhasi, 2008). The expression of osteomyelitis can vary depending on age, nature of the initial infection and immunity of the individual (Pinhasi and Mays, 2008).

Acute infections are usually associated with rapid death rarely affecting the skeleton but it may also stimulate new bone formation (Ortner and Putschard, 1985; Ortner, 2003). Rapid bone formation produces woven bone (active lesions) that typically is the initial stage in many abnormal bone forming lesions caused by infection (Ortner and Putschard, 1985; Ortner, 2003). In chronic or healing stages (healed lesions) the woven bone is remodelled into compact bone (Ortner and Putschard, 1985; Ortner, 2003). However, chronic infectious diseases often have various acute phases. Chronic infections are very informative about the nutritional adequacy of the diet, the state of waste disposal and hygiene in a specific community (Goodman and Martin, 2002). Infectious pathologies, especially when linked with malnutrition, are the largest contributor to morbidity and mortality worldwide (Keusch and Farthing, 1986). The study of nutrition-infection

interactions is important to understand the complexity of the relationships of these factors with immunological status, co-morbidity and mortality (Uljaszek et al., 2012), especially in pre-antibiotic societies.

New bone formation can also be considered an indicator of physiological stress and has been associated with lower socioeconomic status (e.g. Goodman and Martin, 2002; Peck, 2013; Robb et al., 2001), systematic infections (e.g. Goodman and Martin, 2002; Larsen, 2002; Ortner, 2003), malnutrition (e.g. Weston, 2012) and niacin deficiency (Paine and Brenton, 2006), which can leave the individuals more susceptible to pathogens. Deposits of new bone may also be associated with elevated risks of mortality and are therefore informative about ill health (e.g. DeWitte and Wood, 2008).

## 1.3. Stable isotope analysis

Analysis of stable isotope ratios from mineralized tissue has been widely used for dietary reconstruction. This technique is based on the assumption that “you are what you eat (plus a few ‰)” (DeNiro and Epstein, 1976), as a consumer's tissues reflect the isotopic array of the ingested foods.

There is enrichment in  $\delta^{13}\text{C}$  in an animal's body tissues relative to its diet due to the fractionation that occurs during the tissue's formation (Van der Merwe and Vogel, 1978). Consumers have a carbon fractionation factor (enrichment in  $\delta^{13}\text{C}$ ) of approximately 5‰ in their bone collagen relative to their diet (Ambrose and Norr, 1993; Van der Merwe and Vogel, 1978) and an enrichment of 1‰ between trophic levels (DeNiro and Epstein, 1978; Tieszen et al., 1983). There is an increment in  $\delta^{15}\text{N}$  of 3‰–5‰ between trophic levels when compared with consumer's diet (Bocherens and Drucker, 2003; Minagawa and Wada, 1984; Schoeninger and DeNiro, 1984; Schoeninger et al., 1983). This fractionation enables the use of stable nitrogen isotopes ( $\delta^{15}\text{N}$ ) to infer trophic level and high  $\delta^{15}\text{N}$  recorded in bone collagen usually indicates high-protein diets (Sponheimer et al., 2003). There are other factors that can raise bone  $\delta^{15}\text{N}$ , such as aridity (Ambrose and DeNiro, 1986; Heaton, 1987; Heaton et al., 1986; Sealy et al., 1987), physiological (Deschner et al., 2012; D'Ortenzio et al., 2015; Gaye-Siessegger et al., 2004; Katzenberg and Lovell, 1999; Oelbermann and Scheu, 2001) or protein stress (Hobson et al., 1993; Steele and Daniel, 1978).

Previous research on archaeological samples with and without lesions indicative of leprosy showed no significant differences in  $\delta^{13}\text{C}$  or  $\delta^{15}\text{N}$ , suggesting that there were no dietary differences between the two groups (Bayliss et al., 2004; Linderholm and Kjellström, 2011). However, other studies showed marked differences between individuals who survived childhood and those who did not (Beaumont et al., 2015; Reitsema et al., 2016), with the ones who survived having higher animal protein in their post-weaning diets (Reitsema et al., 2016) suggesting that investigation of dietary protein, using stable isotopic analysis, might be used to better understand disease and physiological stress in past populations. Skeletal indicators of physiological stress, such as low stature and *cribra orbitalia*, have also been related to long-term effects on health throughout reduced lifespan (Watts, 2013) and increased risk of death during epidemics (DeWitte and Hughes-Morey, 2012; DeWitte and Wood, 2008).

## 1.4. Diet at Tomar

People living in Tomar had a complex diet, low in terrestrial animal protein and high in aquatic protein intake, despite its inland location (Curto et al., 2018). Being controlled by religious military orders (Conde, 1996; Valente, 1998), it is possible that their presence in the town would have an impact on the general population particularly on their diet (Curto et al., 2018), due to religious fasting (Barber and Bate, 2002; Müldner et al., 2009; Müldner and Richards, 2007; Salamon et al., 2008). Fish was an expensive food source, particularly further away from the coast (Gonçalves, 2004; Vicente, 2013), therefore higher amounts of fish consumption may reflect higher socio-economic status

(Curto et al., 2018).

There were no significant differences found between sexes or age groups for bone collagen  $\delta^{13}\text{C}$  and  $\delta^{34}\text{S}$ , however  $\delta^{15}\text{N}$  did differ significantly with age (lower  $\delta^{15}\text{N}$  in older individuals), which may be related to tooth loss in old individuals (Curto et al., 2018). There was one outlier, a young adult male, with higher values of both  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  and lower  $\delta^{34}\text{S}$  than the other skeletons analysed, suggesting he may be an outsider (Curto et al., 2018). There were no differences between inferred social status, estimated through burial type and proximity to the church (Curto et al., 2018).

### 1.5. Research questions and predictions

The main objective of this study is to determine if there is a link between diet and health assessed by  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  ratios from bone collagen in skeletons that retain evidence of non-specific disease. The stable isotope ratios from long bones' collagen are a long-term measure of dietary protein consumed by an individual over a period of about 10 years of life (Hedges et al., 2007). Thus, we seek to determine if longer term diet corresponds with disease at the point of death. Our predictions are as follows:

Protein malnutrition over a long period of time impairs the immune system and increases the likelihood of an individual contracting an infectious disease (e.g. Calder, 2013; Scrimshaw and SanGiovanni, 1997; Woodward, 1998; Calder and Jackson, 2000; Woodward, 2001). Therefore, individuals with skeletal signs of infectious diseases might have had different diets than those without skeletal lesions. Skeletons with signs of infection might have had a diet poorer in animal protein, than the individuals without lesions, which might have lowered their resistance to disease (e.g. Calder, 2013; Kuvibidila et al., 1993; Scrimshaw and SanGiovanni, 1997; Woodward, 1998; Calder and Jackson, 2000; Woodward, 2001; Ulijaszek et al., 2012; Weston, 2012).

$\delta^{15}\text{N}$  in particular are very informative of trophic level and high  $\delta^{15}\text{N}$  usually indicate high-protein diets (Schoeninger et al., 1983; Minagawa and Wada, 1984; Schoeninger and DeNiro, 1984; Bocherens and Drucker, 2003). Therefore we predict that skeletons without signs of infectious disease have higher  $\delta^{15}\text{N}$  than the ones with skeletal lesions. However, there are other factors that can raise the  $\delta^{15}\text{N}$  including physiological (Katzenberg and Lovell, 1999; Oelbermann and Scheu, 2001; Gaye-Siessegger et al., 2004; Deschner et al., 2012; D'Ortenzio et al., 2015) and/or nutritional stress (Steele and Daniel, 1978; Hobson et al., 1993), which have been associated with  $\delta^{15}\text{N}$  increase due to protein catabolism. In prolonged cases of disease, nutritional or physiological stress, dietary protein cannot adequately replace nitrogen losses (Grossman et al., 1945; Powanda, 1977; Welle, 1999). Consequently, the body proteins are recycled resulting in enriched  $\delta^{15}\text{N}$  (e.g. Steele and Daniel, 1978; Hobson et al., 1993; Deschner et al., 2012; D'Ortenzio et al., 2015).

Periostitis generally reflects a reaction to pathologic changes of the underlying bone, or part of it, but can also result from trauma and/or inflammation of the surrounding tissues (Ortner and Putschard, 1985; Ortner, 2003). Generalised infections (various bones with periostitis and/or osteomyelitis), on the other hand, might represent severe infections which spread across the body (Ortner and Putschard, 1985; Ortner, 2003). However, the presence of skeletal lesions can also represent good physiological state, allowing these individuals to survive long enough to the disease for it to be visible on their bones (Wood et al., 1992). Periostitis reflects physiological stress and morbidity but frequently represents later phases of the inflammation and succeeding recovery from the stress incident (Klaus, 2014). For this reason bone collagen  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  from skeletons without lesions (and other skeletal markers of physiological stress; Curto et al., 2018) will be compared with bone collagen  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  from 1) skeletons with only healed tibial periostitis, 2) skeletons with non-specific generalised infections and 3) skeletons with venereal syphilis.

Woven bone is produced during rapid bone formation and when it is

observed in adults it is considered of pathological origin (Ortner and Putschard, 1985; Ortner, 2003). Since in chronic or healing stages the woven bone is rapidly remodelled into compact bone, woven bone is considered a lesion which was active *perimortem*, while compact bone is considered a lesion which was healed *perimortem* (Ortner and Putschard, 1985; Ortner, 2003). Chronic infectious diseases can also have various acute phases and be very informative about the nutritional adequacy of the diet in a specific community (Goodman and Martin, 2002). Therefore, bone collagen  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  from skeletons without lesions (and other skeletal markers of physiological stress) will be compared with bone collagen  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  from 1) skeletons with only active lesions, 2) skeletons with only healed lesions and 3) skeletons with both healed and active lesions. Since Protein malnutrition impairs the immune system (e.g. Calder, 2013; Scrimshaw and SanGiovanni, 1997; Woodward, 1998; Calder and Jackson, 2000; Woodward, 2001), we predict that skeletons without lesions have higher  $\delta^{15}\text{N}$  than those with lesions, with the ones with only active lesions having the lowest  $\delta^{15}\text{N}$ . The skeletons with only healed lesions are expected to have  $\delta^{15}\text{N}$  similar to the skeletons without lesions as they survived the disease long enough for the bone to remodel into compact bone (Ortner and Putschard, 1985; Ortner, 2003; Wood et al., 1992).

## 2. Materials and methods

Santa Maria do Olival necropolis, at Tomar (Fig. 1), is one of the largest in Europe (6792 individuals recovered: 4991 adults and 1801 non-adults) but has not been continuously studied yet. Even though Tomar was a Templar town the distribution of the skeletons, of all ages and both sexes, within the necropolis suggests that Santa Maria do




Fig. 1. Map of Portugal showing the location of Tomar. Adapted from d-maps.com.

**Table 1**  
Summary of type and number of samples collected.

Adults sampled	Non-specific						Specific		
	Localised			Generalised			Venereal syphilis		
	Active	Healed	Active & Healed	Active	Healed	Active & Healed	Active	Healed	Active & Healed
Young	0	0	2	2	0	0	0	0	1
Mature	2	4	0	0	0	1	0	0	1
Old	1	0	1	0	0	2	0	0	0
Undetermined	1	3	2	0	0	0	0	0	0
Total	4	7	5	2	0	3	0	0	2
	16			5			2		
	21								

Olival collection represents the general population of Tomar and not, or at least not only, the individuals from the military orders (Curto et al., 2018).

Bone collagen stable isotope data (carbon, nitrogen and sulphur) from 32 human adult tibiae (15 females; 18 males) and 13 faunal remains (2 wild *Sus*; 2 domestic *Sus*; 1 juvenile *Sus*; 1 *Canidae*; 3 *Bos*; 1 *Equus*; 3 *Ovicapridae*) from Tomar (11th – 17th century) were previously analysed to reconstruct the general diet of the population (Curto et al., 2018). These are reused here and compared to new isotope data from skeletons with signs of disease (Table 1). These data are compared to new isotope ratios from 23 adult individuals (8 females; 14 males; 1 undetermined) with skeletal lesions compatible with non-specific (n = 21) and specific (venereal syphilis, n = 2) infectious diseases.

All samples are from Santa Maria do Olival graveyard (areas 13 to 20; 11th to 17th centuries) in Tomar. The individuals without lesions (n = 32), previously analysed (Curto et al., 2018), were used to estimate the baseline diet at Tomar and were selected based on the absence of skeletal lesions or skeletal stress markers (see Curto et al., 2018 for more detail; the outlier was not considered for this study). There were no significant differences found between sexes or inferred social status, estimated through burial type and proximity to the church (Curto et al., 2018).

### 2.1. Estimating age and sex

Sex was estimated based on pelvic (Phenice, 1969; Buikstra and Ubelaker, 1994) and cranial features (Buikstra and Ubelaker, 1994). Adult age at death estimates employed a combination of skeleton maturation (Scheuer and Black, 2000), pubic symphysis degeneration (Brooks and Suchey, 1990; Buikstra and Ubelaker, 1994) and auricular surface degeneration (Lovejoy et al., 1985). The skeletons analysed were grouped as young (18–30 years; n = 5), mature (31–60 years; n = 8) and old (60 + years; n = 4) adults; for six skeletons it was not possible to estimate age.

### 2.2. Signs of infection

From the 23 skeletons with lesions (Table 1), 21 have signs of non-specific infectious diseases and 2 have lesions compatible with specific infections (venereal syphilis). The 23 individuals were grouped in two different ways: a) active (n = 6), healed (n = 7) and a combination of both active and healed lesions (n = 10); b) Skeletons with only healed tibial periostitis (n = 7), those with non-specific (n = 5) and specific (n = 2) infectious diseases, while individuals who did not fit into these groups (n = 9) were not considered for this analysis. Figs. 2–4 show examples of the different lesion stages analysed.

Skeletal lesions were considered to be from possible infectious causes if abnormal bone formation or bone formation and destruction, compatible with periostitis or osteomyelitis (Ortner and Putschard, 1985; Buikstra and Ubelaker, 1994; Aufderheide and Rodríguez-Martín,

1998; Ortner, 2003), were present and not associated with trauma. Periostitis usually represents pathologic changes resulting in new bone growth, which is remodelled into lamellar bone during the healing process, but it can also result from inflammation of the surrounding tissues following a trauma (Ortner and Putschard, 1985; Ortner, 2003).

For this study, lesions scored 2 (markedly accentuated longitudinal striations on the surface of cortical bone; Steckel et al., 2006) to 5 (extensive periosteal reaction involving over half of the diaphysis, with cortical expansion, pronounced deformation; Steckel et al., 2006) were considered periostitis. Lesions that were scored as 6 (involving most of the diaphysis with cloacae; Steckel et al., 2006) were taken as evidence of osteomyelitis. Periostitis or osteomyelitis associated with fractures was not considered for this study.

Lesions with unremodelled woven bone were considered active at the time of death (Ortner and Putschard, 1985; Ortner, 2003). Rapidly formed woven bone is poorly organized and has a porous appearance due to the loose organization of the mineralized osteoid fibres (Ortner and Putschard, 1985; Ortner, 2003). Markedly accentuated longitudinal striations and compact bony growth, without the presence of woven bone, were considered healed lesions (Ortner and Putschard, 1985; Ortner, 2003). The presence of both compact bony growth and woven bone was considered a combination of both healed and active lesions. The skeletons with only active lesions represent infectious diseases active *perimortem* and the ones with only healed lesions represent healed individuals. Skeletons with a combination of both types of lesions represent chronic infections, to which the individuals survived long enough to the disease for the bone to heal but with the disease still present. The skeletons with the different lesions (healed, active and both) were combined and compared with the individuals without lesions, by age group: young without lesions (n = 8); young with lesions (n = 5); mature without lesions (n = 13); mature with lesions (n = 8); old without lesions (n = 4) and old with lesions (n = 4).

Since tibial periostitis is frequently used as an indicator of physiological stress (e.g. DeWitte, 2010; Robb et al., 2001) and can be caused by a variety of factors, including trauma, only individuals with bilateral healed periostitis on the tibiae were selected (markedly accentuated longitudinal striations; score 2; Steckel et al., 2006). The cases of venereal syphilis were diagnosed due to the presence of *caries sicca*, a sign specifically characteristic of venereal syphilis (Ortner and Putschard, 1985; Aufderheide and Rodríguez-Martín, 1998; Ortner, 2003). These groups with signs of infections were then compared with the skeletons without lesions (n = 32; Curto et al., 2018).

The skeletons were grouped in different ways to better understand how diet may affect the susceptibility to generalised infections (by grouping non-specific generalised infections, specific generalised infections and individuals with only healed tibial periostitis) or the ability to recover from infectious diseases (by grouping the skeletons as having active, healed or a combination of both active and healed lesions).

Only tibiae collagen was analysed in an attempt to estimate the average long term diet of the individuals and avoid stable isotopes data



Fig. 2. Example of healed tibial periostitis (skeleton 15.96).

that may represent different diet and/or metabolism during the disease. Following the attempt to avoid stable isotope values related to faster bone remodelling and therefore more recent diet, samples were only collected at areas of the bone without any sign of lesions.

### 2.3. Collagen extraction and analysis

Collagen extraction was done following Longin (1971), Brown et al. (1988) and Richards and Hedges (1999). The collagen samples were weighed into tin capsules and combusted into CO<sub>2</sub> and N<sub>2</sub> in an isotope-ratio mass spectrometer at NERC Isotope Geosciences Facility and HERCULES laboratory. At NERC,  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  were calibrated using an in-house reference material M1360p (powdered gelatine from British Drug Houses) with expected  $\delta$  values of  $-20.32\text{‰}$  (calibrated against CH<sub>7</sub>, IAEA) and  $+8.12\text{‰}$  (calibrated against N-1 and N-2, IAEA) for carbon and nitrogen respectively. Samples were run in duplicate and the  $1\sigma$  reproducibility for mass spectrometry controls for these analyses were  $\delta^{15}\text{N} = \pm 0.08\text{‰}$  and  $\delta^{13}\text{C} = \pm 0.07\text{‰}$ . At HERCULES Laboratory,  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  were calibrated using IAEA-CH-6 (sucrose,  $-10.449\text{‰}$ ), IAEA-CH-7 (polyethylene,  $-32.151\text{‰}$ ), IAEA-N-1 (ammonium sulphate,  $+0.4\text{‰}$ ) and IAEA-N-2 (ammonium sulphate,  $+20.3\text{‰}$ ). Measurement errors were less than  $\pm 0.1\text{‰}$  for  $\delta^{13}\text{C}$  and  $\pm 0.2\text{‰}$  for  $\delta^{15}\text{N}$ .

Mann-Whitney U non-parametric tests were used for pair-wise comparisons and Kruskal-Wallis non-parametric tests were used to compare more than two groups. All statistics were computed in SPSS 24 for Windows and  $p$ -values  $\leq 0.05$  were considered statistically significant.

## 3. Results

### 3.1. Bone collagen $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ of skeletons with generalised infections or healed tibial periostitis compared to skeletons without lesions

Osteomyelitis was only observed in the skeletons with venereal syphilis (skeletons 16.225 and 18.158) and skeleton 16.255

( $\delta^{13}\text{C} = -18.7\text{‰}$ ;  $\delta^{15}\text{N} = 10.0\text{‰}$ ), a mature male with osteomyelitis on the right tibia. Therefore, the results from this study are focused mainly on lesions within the scope of periostitis.

Fig. 5 illustrates the  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  for skeletons without lesions ( $n = 32$ ; Curto et al., 2018), with only healed tibial periostitis ( $n = 7$ ) and those with generalised specific ( $n = 2$ ) and non-specific ( $n = 5$ ) infections. There is one outlier with healed tibial periostitis ( $\delta^{13}\text{C} = -15.6\text{‰}$ ;  $\delta^{15}\text{N} = 11.5\text{‰}$ ) that seems to have very different diet from the general population and therefore was not considered for the statistical analysis. Among the individuals with skeletal lesions, the ones with healed tibial periostitis ( $n = 6$ ; one is an outlier) have the highest mean values for both  $\delta^{13}\text{C}$  ( $-18.0 \pm 1.1\text{‰}$ ; Table 2) and  $\delta^{15}\text{N}$  ( $10.9 \pm 0.7\text{‰}$ ; Table 2), while those with non-specific generalised infections ( $n = 5$ ) have the lowest mean for  $\delta^{13}\text{C}$  ( $-18.7 \pm 0.8\text{‰}$ ; Table 1) and  $\delta^{15}\text{N}$  ( $9.9 \pm 0.4\text{‰}$ ; Table 1). The skeletons with venereal syphilis ( $n = 2$ ) have similar mean values ( $\delta^{13}\text{C} = -18.5 \pm 0.2\text{‰}$ ;  $\delta^{15}\text{N} = 11.2 \pm 0.3\text{‰}$ ) to the skeletons without lesions ( $n = 32$ ;  $\delta^{13}\text{C} = -18.6 \pm 0.5\text{‰}$ ;  $\delta^{15}\text{N} = 10.8 \pm 0.8\text{‰}$ ) and those with only healed tibial periostitis ( $n = 6$ ), however the sample size is too small for an appropriate statistical analysis. The difference in  $\delta^{15}\text{N}$  between skeletons with non-specific generalised infection ( $\delta^{13}\text{C} = -18.7 \pm 0.8\text{‰}$ ;  $\delta^{15}\text{N} = 9.9 \pm 0.4\text{‰}$ ) and healed periostitis ( $\delta^{13}\text{C} = -18.1 \pm 1.2\text{‰}$ ;  $\delta^{15}\text{N} = 11.2 \pm 0.4\text{‰}$ ) is highly significant ( $p < 0.003$ ; Table 2) as is the difference between skeletons with non-specific generalised infection and those without lesions ( $\delta^{13}\text{C} = -18.5 \pm 0.7\text{‰}$ ;  $\delta^{15}\text{N} = 10.9 \pm 0.9\text{‰}$ ) ( $p < 0.004$ ; Table 1). There are no statistically significant differences for  $\delta^{13}\text{C}$  ( $p > 0.53$ ; Table 2) or between skeletons without lesions and skeletons with only healed tibial periostitis for both  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  ( $p > 0.20$ ; Table 2).

### 3.2. Bone collagen $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ of skeletons with lesions compared to skeletons without lesions, by age groups

Fig. 6 illustrates  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  for individuals with (including healed, active or a combination of both lesions) and without lesions by age group



Fig. 3. Example of a lesion combining active and healed periosteal reactions (skeleton v5.22).



Fig. 4. Example of healed osteomyelitis from an individual with syphilis (skeleton 20.240).




Fig. 5.  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  (‰) for individuals without lesions, with only healed periostitis, with non-specific generalised infections and with treponematosi. Data from skeletons without lesions previously analysed in Curto et al. (2018).

(Table 3). Young adults without lesions ( $n = 8$ ) have higher  $\delta^{13}\text{C}$  ( $-18.5 \pm 0.4\text{‰}$ ) and  $\delta^{15}\text{N}$  ( $11.4 \pm 0.7\text{‰}$ ) than the ones with lesions ( $n = 5$ ;  $\delta^{13}\text{C} = -18.8 \pm 0.4\text{‰}$ ;  $\delta^{15}\text{N} = 10.5 \pm 0.8\text{‰}$ ) but still falling within the two standard deviations of each other and the general sample without lesions. There is no statistically significant differences in  $\delta^{13}\text{C}$  or  $\delta^{15}\text{N}$  for the mature (without lesions:  $n = 13$ ;  $\delta^{13}\text{C} = -18.6 \pm 0.6\text{‰}$ ;  $\delta^{15}\text{N} = 10.5 \pm 0.7\text{‰}$ ; with lesions:  $n = 8$ ;  $\delta^{13}\text{C} = -18.5 \pm 0.5\text{‰}$ ;

Table 2

Mean, standard deviation and non parametric tests for  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  (‰) of individuals without lesions, with healed periostitis and with generalised infections (without outliers). Data from skeletons without lesions previously analysed in Curto et al. (2018).

	N	Mean $\pm$ sd		Non parametric test		
		$\delta^{13}\text{C}$	$\delta^{15}\text{N}$	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$	
Healthy	32	$-18.5 \pm 0.7$	$10.9 \pm 0.9$	95.00	80.00	Mann-Whitney U sig
Healed periostitis	6	$-18.0 \pm 1.1$	$10.9 \pm 0.6$	0.49	0.21	
Healthy	32	$-18.5 \pm 0.7$	$10.9 \pm 0.9$	74.00	19.00	Mann-Whitney U sig
Non-specific generalised infection	5	$-18.7 \pm 0.8$	$9.9 \pm 0.4$	0.74	0.00	
Healed periostitis	6	$-18.0 \pm 1.1$	$10.9 \pm 0.6$	20.00	7.00	Mann-Whitney U sig
Non-specific generalised infection	5	$-18.7 \pm 0.8$	$9.9 \pm 0.4$	0.53	0.00	

$\delta^{15}\text{N} = 10.7 \pm 0.7\text{‰}$ ) and old adults (without lesions:  $n = 4$ ;  $\delta^{13}\text{C} = -18.6 \pm 0.3\text{‰}$ ;  $\delta^{15}\text{N} = 10.7 \pm 1.2\text{‰}$ ; with lesions:  $n = 4$ ;  $\delta^{13}\text{C} = -18.4 \pm 0.3\text{‰}$ ;  $\delta^{15}\text{N} = 10.3 \pm 0.4\text{‰}$ ) ( $p > 0.38$ ; Table 3).

### 3.3. Bone collagen $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ of skeletons with active, healed or a combination of both lesions compared to skeletons without lesions

The only healed lesions were found within the mature adults group (Fig. 6). Results show there is no statistically significant difference in  $\delta^{13}\text{C}$  or  $\delta^{15}\text{N}$  when the skeletons without visible lesions ( $n = 32$ ;  $\delta^{13}\text{C} = -18.6 \pm 0.5\text{‰}$ ;  $\delta^{15}\text{N} = 10.8 \pm 0.8\text{‰}$ ; Table 4) were compared with the skeletons with healed ( $n = 6$ ;  $\delta^{13}\text{C} = -18.4 \pm 0.4\text{‰}$ ;  $\delta^{15}\text{N} = 10.8 \pm 0.7\text{‰}$ ;  $p = 0.53$ ; Table 4), active ( $n = 6$ ;  $\delta^{13}\text{C} = -18.5 \pm 0.7\text{‰}$ ;  $\delta^{15}\text{N} = 10.5 \pm 0.7\text{‰}$ ;  $p = 0.72$ ; Table 4) or a combination of both lesions ( $n = 10$ ;  $\delta^{13}\text{C} = -18.4 \pm 0.2\text{‰}$ ;  $\delta^{15}\text{N} = 10.7 \pm 0.8\text{‰}$ ;  $p = 0.24$ ; Table 4).

## 4. Discussion

### 4.1. Bone collagen $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ of skeletons with generalised infections or healed tibial periostitis compared to skeletons without lesions

The  $\delta^{15}\text{N}$  enrichment observed in skeletons with only healed tibial periostitis ( $N = 6$ , without the outlier), when compared to those with non-specific generalised infections ( $n = 5$ ), may represent evidence of chronic physiological stress (Steele & Daniel, 1978; Hobson et al., 1993; Gaye-Siessegger et al., 2004; Fuller et al., 2005; Deschner et al., 2012; D'Ortenzio et al., 2015; Scorrano et al., 2014). However, the individuals with non-specific generalised infections ( $n = 5$ ) were also exposed to chronic physiological stress and survived long enough for it to be observable in their bones (Wood et al., 1992); yet they display lower  $\delta^{15}\text{N}$  ( $9.9 \pm 0.4\text{‰}$ ) than the individuals without lesions ( $n = 32$ ;  $\delta^{15}\text{N} = 10.8 \pm 0.8\text{‰}$ ), those with only healed tibial periostitis ( $n = 6$ ;  $\delta^{15}\text{N} = 10.9 \pm 0.7\text{‰}$ ) and the ones with venereal syphilis ( $n = 2$ ;  $\delta^{15}\text{N} = 10.5 \pm 0.6\text{‰}$ ).

The only skeleton with osteomyelitis (16.255), besides the ones with




Fig. 6.  $\delta^{13}C$  and  $\delta^{15}N$  (‰) for individuals with and without lesions, by age group (means calculated without outliers). Data from skeletons without lesions previously analysed in Curto et al. (2018).

Table 3

Mean, standard deviation and non parametric tests for  $\delta^{13}C$  and  $\delta^{15}N$  (‰) of individuals with and without lesions, by age group. Data from skeletons without lesions previously analysed in Curto et al. (2018).

		N	Mean $\pm$ sd		Non parametric test		
			$\delta^{13}C$	$\delta^{15}N$	$\delta^{13}C$	$\delta^{15}N$	
Young	Without lesions	7	$-18.5 \pm 0.4$	$11.4 \pm 0.7$	7.00	7.00	Mann-Whitney U sig
	With lesions	5	$-18.8 \pm 0.4$	$10.5 \pm 0.8$	0.09	0.09	
Mature	Without lesions	13	$-18.6 \pm 0.6$	$10.5 \pm 0.7$	60.00	59.00	Mann-Whitney U sig
	With lesions	8	$-18.5 \pm 0.5$	$10.7 \pm 0.7$	0.51	0.49	
Old	Without lesions	4	$-18.6 \pm 0.3$	$10.7 \pm 1.2$	5.00	6.00	Mann-Whitney U sig
	With lesions	4	$-18.4 \pm 0.3$	$10.3 \pm 0.4$	0.39	0.56	

venereal syphilis, has similar  $\delta^{13}C$  ( $-18.7\text{‰}$ ) and  $\delta^{15}N$  ( $10.0\text{‰}$ ) to the individuals with non-specific generalised infections ( $\delta^{13}C = -18.7 \pm 0.8\text{‰}$ ;  $\delta^{15}N = 9.9 \pm 0.4\text{‰}$ ; Table 2), suggesting that a diet lower in animal protein might have made him more susceptible to infectious disease (e.g. Kuvibidila et al., 1993; Scrimshaw and SanGiovanni, 1997; Woodward, 1998; Calder and Jackson, 2000; Woodward, 2001). Venereal syphilis is a sexually transmitted disease and human hosts have no natural immunity to pathogenic treponemes (Kiple, 1993). Therefore, the immune system of the individuals before the disease is not as relevant to the individuals' susceptibility to these infections. However, good health prior to venereal syphilis infection may prolong the individual's survival (not only to the treponeme but also to other infections through skin ulcers which increase exposure to other pathogens) and increase the amount and severity of the lesions (Wood et al., 1992).

The skeletons without lesions were also carefully chosen not only based on the absence of infectious lesions (including tibial periostitis)

but also other physiological stress indicators such as cribra orbitalia, porotic hyperostosis, enamel hypoplasias and stature above the average for the population under study (Curto et al., 2018). Even so, the skeletons with only healed tibial periostitis have similar  $\delta^{13}C$  and  $\delta^{15}N$  to those without any sign of physiological stress (Fig. 5).

The osteological paradox (Wood et al., 1992) may explain the higher  $\delta^{13}C$  and  $\delta^{15}N$  for the skeletons with only healed tibial periostitis when compared to the ones with non-specific generalised infections (Fig. 5 & Table 2). It is possible that the skeletons with only healed tibial periostitis had a diet richer in animal protein and therefore were more resistant to diseases (e.g. Calder, 2013; Kuvibidila et al., 1993; Scrimshaw and SanGiovanni, 1997; Woodward, 1998; Calder and Jackson, 2000; Woodward, 2001; Ulijaszek et al., 2012; Weston, 2012) than those who had non-specific generalised infections. It has been argued that individuals with healed periostitis are of lower frailty, having a lower risk of death (e.g. DeWitte, 2010; Ortner, 2003; Wood et al., 1992).

**Table 4**

Mean, standard deviation and non parametric tests for  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  (‰) of individuals with different types of lesions and without lesions (without outliers). Data from skeletons without lesions previously analysed in Curto et al. (2018).

	N	Mean $\pm$ sd		Non parametric test		
		$\delta^{13}\text{C}$	$\delta^{15}\text{N}$	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$	
Without lesions	32	$-18.6 \pm 0.5$	$10.8 \pm 0.8$	66.00	77.00	Mann-Whitney U
Healed lesions	6	$-18.4 \pm 0.4$	$10.8 \pm 0.7$	0.53	0.89	sig
Without lesions	32	$-18.6 \pm 0.5$	$10.8 \pm 0.8$	87.00	73.00	Mann-Whitney U
Active lesions	6	$-18.5 \pm 0.7$	$10.5 \pm 0.7$	0.72	0.36	sig
Without lesions	32	$-18.6 \pm 0.5$	$10.8 \pm 0.8$	120.00	134.00	Mann-Whitney U
Active & healed lesions	10	$-18.4 \pm 0.2$	$10.7 \pm 0.8$	0.24	0.44	sig

The diet of the population under study was complex and likely included food sources from outside Tomar (Curto et al., 2018). The diet of these individuals was poor in terrestrial protein and rich in aquatic protein ( $\delta^{13}\text{C} = -18.6\text{‰}$ ;  $\delta^{15}\text{N} = 10.8\text{‰}$ ;  $\delta^{34}\text{S} = 13.1\text{‰}$ ; Curto et al., 2018). Stable isotope values are similar for males and females but the young adults have higher  $\delta^{15}\text{N}$  ( $11.4 \pm 0.6\text{‰}$ ) than the old adults ( $10.6 \pm 0.8\text{‰}$ ), suggesting a higher animal protein intake for the young individuals (Curto et al., 2018). The high  $\delta^{15}\text{N}$  from skeletons without lesions seem to be related with higher aquatic protein intake (Curto et al., 2018), which may be related with these individuals having better health than those with signs of infection. Since fish was expensive (Gonçalves, 2004) and the military orders had angling rights (Vicente, 2013) it is also possible that the individuals without skeletal stress markers, or only healed tibial periostitis, had a higher socioeconomic status. Socioeconomic status may also have an impact on an individual's diet, not only directly on their diet but also the type of pathogens they would be exposed to.

The effect of protein malnutrition on the immune system is well known (Calder, 2013; Kuvibidila et al., 1993; Scrimshaw and SanGiovanni, 1997; Woodward, 1998, 2001; Calder and Jackson, 2000) and the possibility of dietary differences being present before the disease cannot be excluded.  $\delta^{15}\text{N}$  were significantly different between skeletons with non-specific generalised infections and those without lesions ( $p < 0.004$ ) or with only healed tibial periostitis ( $p < 0.003$ ). The higher  $\delta^{15}\text{N}$  observed in the two individuals with venereal syphilis, may not be related to physiological stress but may be due to the nature of the disease instead (sexually transmitted infection) and the  $\delta^{15}\text{N}$  might suggest a richer diet that could have allowed survival despite the disease and susceptibility to other pathogens. The possibility of these  $\delta^{15}\text{N}$  differences being related with social status cannot be excluded. Various studies suggest dietary differences between sex and social status in Medieval times (e.g. Adamson, 2004; Kjellström et al., 2009; Linderholm et al., 2008; Polet and Katzenberg, 2003; Schutkowski et al., 1999; Reitsema et al., 2010; Reitsema and Vercellotti, 2012). However, a previous study showed no significant stable isotope data between individuals of different sex or social status in Tomar (Curto et al., 2018).

There are two outliers among the skeletons sampled for isotopic analysis (Fig. 5), one without lesions and another one with healed tibial periostitis. The skeleton without lesions, a young adult male, might be an outsider as his sulphur isotopes ratios (9.3‰) differ from the other individuals without lesions (mean  $\delta^{34}\text{S} = 13.1\text{‰}$ ; Curto et al., 2018). This skeleton was not considered for the statistical analysis. There are no sulphur isotopes values for the outlier with healed tibial periostitis but  $\delta^{13}\text{C}$  ( $-15.6\text{‰}$ ) and  $\delta^{15}\text{N}$  (11.5‰) are similar to those of the outlier without lesions ( $\delta^{13}\text{C} = -15.4\text{‰}$ ;  $\delta^{15}\text{N} = 12.3\text{‰}$ ).

#### 4.2. Bone collagen $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ of skeletons with lesions compared to skeletons without lesions

The values for the young adults show a statistical trend towards a significance ( $p < 0.09$ ; Table 3) difference in both  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  between skeletons with ( $n = 5$ ) and without ( $n = 8$ ) lesions. Young individuals without lesions have higher  $\delta^{13}\text{C}$  ( $-18.5 \pm 0.4\text{‰}$ ) and  $\delta^{15}\text{N}$  ( $11.4 \pm 0.7\text{‰}$ ) than those with lesions ( $\delta^{13}\text{C} = -18.8 \pm 0.4\text{‰}$ ;  $\delta^{15}\text{N} = 10.5 \pm 0.8\text{‰}$ ), which may suggest that the individuals with lesions may have had a diet with lower animal protein (Fig. 6). There is no difference for mature ( $p > 0.49$ ; Table 3) and old ( $p > 0.39$ ; Table 3) individuals with or without lesions. Previous research on archaeological samples showed marked differences between individuals who survived childhood and those who did not (Beaumont et al., 2015; Reitsema et al., 2016), with the ones who survived having higher animal protein in their post-weaning diets (Reitsema et al., 2016) suggesting that diet at younger ages can have a high impact on the health status of an individual. The impact of diet on an individual's health might be prolonged throughout adult life as well. The young adult skeletons analysed do not have healed lesions, only active or a combination of both active and healed lesions, meaning that they died during acute phases of the disease (Ortner and Putschard, 1985; Ortner, 2003; Turner-Walker, 2008).

#### 4.3. Bone collagen $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ of skeletons with active, healed or a combination of both lesions compared to skeletons without lesions

The absence of significant differences in  $\delta^{13}\text{C}$  or  $\delta^{15}\text{N}$  between individuals without lesions ( $n = 32$ ;  $\delta^{13}\text{C} = -18.6 \pm 0.5\text{‰}$ ;  $\delta^{15}\text{N} = 10.8 \pm 0.8\text{‰}$ ; Table 4) and those with healed ( $n = 6$ ;  $\delta^{13}\text{C} = -18.4 \pm 0.4\text{‰}$ ;  $\delta^{15}\text{N} = 10.8 \pm 0.7\text{‰}$ ; Table 4), active ( $n = 6$ ;  $\delta^{13}\text{C} = -18.5 \pm 0.7\text{‰}$ ;  $\delta^{15}\text{N} = 10.5 \pm 0.7\text{‰}$ ; Table 4) or a combination of both lesions ( $n = 10$ ;  $\delta^{13}\text{C} = -18.4 \pm 0.2\text{‰}$ ;  $\delta^{15}\text{N} = 10.7 \pm 0.8\text{‰}$ ;  $p = 0.24$ ; Table 4) suggests that diet may have a higher impact on the susceptibility to chronic generalised infections than to infectious disease in general. It is therefore important to take into account the severity and stage of the disease. The  $\delta^{15}\text{N}$  average is slightly higher for the individuals without lesions (10.8‰;  $n = 32$ ) than for the one ones with active lesions (10.5‰;  $n = 6$ ; Table 4). This slight difference may indicate that the individuals without lesions had a diet richer in animal protein than those with active lesions, however the sample size is too small to make conclusions.

### 5. Study limitations

One of the limitations of this study is the impossibility of knowing the cause of death for the individuals analysed, alongside it not being possible to know which diseases caused most of the lesions and how long the individuals survived with the infections. The presence of skeletal lesions can represent an adaptation to a pathological condition (Ortner, 2003) indicating that the individual survived long enough for evidence to manifest in the skeletal tissues (Wood et al., 1992). The absence of skeletal lesions is ambiguous; it can indicate either good health, or a fast death as result of an acute disease (DeWitte and Stojanowski, 2015; Siek, 2013; Ortner, 2003; Wood et al., 1992). Another limitation is that, while individuals with poorer nutrition are less resistant to infectious diseases, infectious disease further lowers nutritional status (e.g. Mata et al., 1971; Martorell et al., 1980; Calder, 2013; Scrimshaw and SanGiovanni, 1997; Calder and Jackson, 2000). Funding limitations, associated with the difficulty of finding skeletons that would fit into the different groups, resulted in small sample sizes that are also a limitation for interpreting the results from this study. Therefore, other researchers may find different results when replicating this study. Future research might explore a multivariate approach to the analyses of isotopic data related to age, sex, burial phase, and burial type. Such an approach may reveal the different ways multiple lines of



evidence can interact to predict isotope data.

### 6. Conclusion

This study is part of a larger project that will compare intra-bone stable isotopic data from sites with and without skeletal lesions compatible with diseases and/or physiological stress. This study explored the dietary differences between individuals with and without skeletal lesions compatible with infectious diseases to better understand the impact of diet on individuals' health status and their susceptibility to infectious disease. There is a highly significant difference in  $\delta^{15}\text{N}$  between skeletons with healed tibial periostitis and non-specific generalised infection, as well as a difference at the margin of statistical significance between skeletons without lesions and those with generalised infections. These results demonstrate that the individuals with non-specific generalised infections had diets lower in animal protein than those without lesions or with only healed tibial periostitis. Poorer

diets may increase susceptibility to pathogens leading more frequently to generalised infections while richer diets might increase the survivorship and ability to heal from infectious diseases. However, the possibility of these isotope ratios being a result of the disease cannot be excluded and more data from different periods of time within the individual's' life is necessary to understand when these differences started to manifest. These results indicate that diet has a higher impact on the health status of young people than mature or old individuals, being linked to selective mortality. Our results demonstrate that while non-specific generalised infections are a sign of ill health and poor diet, only healed tibial periostitis indicate a state of comparatively good overall health and diet.

### Funding

University of Kent 50<sup>th</sup> Anniversary Scholarship; FCT [UID/Multi/04449/2013, IF/01661/2015].

### Appendix

Table A.1

Stable isotope results ( $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ) for the skeletons without lesions, with healed periostitis, unspecific generalised infections, specific generalised infections and localised infections. \*Data from skeletons without lesions previously analysed in Curto et al. (2018).

	Sex	Age	Area	$\delta^{13}\text{C}$	%C	$\delta^{15}\text{N}$	%N	C/N	Type of lesion
*Without lesions	Female	–	13	–18.8	32.1	10.6	11.2	3.4	No lesion
	Female	–	14	–18.6	31.3	12.0	10.4	3.4	No lesion
	Female	Mature	16	–18.3	41.8	10.5	14.8	3.3	No lesion
	Female	Mature	14	–18.9	41.3	10.2	14.6	3.3	No lesion
	Female	Old	14	–18.9	21.9	10.3	7.5	3.4	No lesion
	Female	Old	14	–18.7	41.0	10.5	14.5	3.3	No lesion
	Female	Mature	18	–19.1	40.8	8.9	14.4	3.3	No lesion
	Female	Old	20	–18.7	36.7	9.6	12.9	3.3	No lesion
	Female	Young	20	–18.5	23.2	11.0	7.7	3.4	No lesion
	Female	Old	18	–18.3	34.5	12.3	11.9	3.4	No lesion
	Female	Mature	20	–18.6	23.0	11.0	7.9	3.4	No lesion
	Female	Young	14	–19.0	15.5	11.4	5.2	3.4	No lesion
	Female	Old	14	–18.7	41.7	10.1	14.3	3.4	No lesion
	Female	Young	16	–18.9	18.5	10.6	6.1	3.4	No lesion
	Female	–	19	–17.8	32.7	11.2	11.5	3.3	No lesion
	Male	–	20	–18.8	39.1	10.2	13.7	3.3	No lesion
	Male	–	19	–18.9	41.0	9.6	14.1	3.4	No lesion
	Male	Mature	18	–18.8	36.6	11.2	12.8	3.3	No lesion
	Male	Mature	19	–18.2	35.3	10.4	12.3	3.4	No lesion
	Male	Old	18	–19.0	37.9	10.5	12.5	3.4	No lesion
	Male	Mature	14	–19.4	14.8	11.0	5.2	3.3	No lesion
	Male	–	17	–17.3	43.0	10.4	14.8	3.4	No lesion
	Male	Young	18	–18.1	24.1	12.1	8.4	3.4	No lesion
	Male	Mature	14	–19.1	39.0	9.3	13.3	3.4	No lesion
	Male	Young	14	–17.8	39.2	12.5	14.4	3.2	No lesion
	Male	Young	20	–18.5	40.8	11.7	14.1	3.4	No lesion
	Male	Young	16	–18.3	36.1	10.7	12.6	3.4	No lesion
	Male	Young	14	–15.4	25.7	12.3	9.0	3.3	No lesion
	Male	Mature	14	–17.9	42.1	11.2	14.9	3.3	No lesion
	Male	Mature	15	–18.3	33.7	10.5	11.7	3.4	No lesion
Male	Mature	14	–17.6	32.6	11.6	10.9	3.4	No lesion	
Male	Mature	14	–18.7	41.4	11.8	14.3	3.4	No lesion	
Male	Mature	17	–19.1	22.8	10.7	7.6	3.4	No lesion	
Healed periostitis	–	–	20.596	–17.9	28.3	11.0	9.9	3.3	Healed
	Female	–	14.15	–18.4	32.7	10.1	11.5	3.4	Healed
	Female	Mature	14.392	–19.0	17.5	10.7	5.8	3.5	Healed
	Male	Mature	14.388	–18.3	15.0	11.0	5.0	3.4	Healed
	Male	–	15.191	–15.6	34.0	11.5	11.9	3.3	Healed
	Male	Mature	18.250	–18.6	17.8	11.6	6.0	3.5	Healed
	Male	Mature	3.73	–18.3	15.1	9.6	4.8	3.4	Healed
Unspecific generalised infections	Female	Young	16.169	–18.6	41.9	9.7	15.8	3.3	Active
	Female	Old	14.72	–18.4	28.9	10.1	9.8	3.4	Active & healed
	Female	Old	15.96	–18.4	15.0	10.0	5.0	3.3	Active & healed
	Male	Young	14.21	–19.4	42.9	10.1	16.7	3.3	Active
	Male	Mature	V5.22	–18.5	33.8	9.8	11.6	3.4	Active & healed

(continued on next page)

Table A.1 (continued)

	Sex	Age	Area	$\delta^{13}\text{C}$	%C	$\delta^{15}\text{N}$	%N	C/N	Type of lesion
Specific generalised infections	Male	Young	16.225	−18.6	27.0	11.0	9.2	3.4	Active & healed
	Male	Mature	18.158	−18.3	27.5	11.4	9.5	3.4	Active & healed
Localised lesions	Female	–	18.3	−18.1	15.5	10.0	5.2	3.3	Active
	Female	Young	14.22	−18.9	18.5	10.0	6.1	3.3	Active & healed
	Female	–	17.464	−18.3	34.0	11.9	11.9	3.3	Active & healed
	Male	Mature	16.255	−18.7	34.4	10.0	11.8	3.4	Active
	Male	Mature	18.160	−17.7	17.5	10.9	5.8	3.4	Active
	Male	–	19.42	−18.4	21.4	10.7	7.0	3.3	Active & healed
	Male	Old	14.130	−18.7	21.4	11.0	7.0	3.4	Active
	Male	Old	19.45	−18.0	28.9	10.0	9.8	3.3	Active & healed
	Male	Young	17.556	−18.5	22.1	11.7	7.5	3.4	Active & healed

## References

- Adamson, M.W., 2004. Food in Medieval Times. Greenwood Publishing Group.
- Ambrose, S.H., DeNiro, M.J., 1986. Reconstruction of African human diet using bone collagen carbon and nitrogen isotope ratios. *Nature* 319, 321–324.
- Ambrose, S.H., Norr, L., 1993. Experimental evidence for the relationship of the carbon isotope ratios of whole diet and dietary protein to those of bone collagen and carbonate. In: *Prehistoric Human Bone*. Springer, Berlin, Heidelberg, pp. 1–37.
- Armstrong, G.J., 2003. Bioarchaeology as anthropology. *Archaeological Papers of the American Anthropological Association* 13 (1), 27–40.
- Aufderheide, A.C., Rodriguez-Martin, C., 1998. Human Paleopathology. Cambridge University Press.
- Barber, M., Bate, K., 2002. The Templars: Selected Sources. Manchester University Press.
- Bayliss, A., Popescu, E.S., Beavan-Athfield, N., Ramsey, C.B., Cook, G.T., Locker, A., 2004. The potential significance of dietary offsets for the interpretation of radiocarbon dates: an archaeologically significant example from medieval Norwich. *J. Archaeol. Sci.* 31 (5), 563–575.
- Beaumont, J., Montgomery, J., Buckberry, J., Jay, M., 2015. Infant mortality and isotopic complexity: new approaches to stress, maternal health, and weaning. *Am. J. Phys. Anthropol.* 157 (3), 441–457.
- Bocherens, H., Drucker, D., 2003. Trophic level isotopic enrichment of carbon and nitrogen in bone collagen: case studies from recent and ancient terrestrial ecosystems. *Int. J. Osteoarchaeol.* 13 (1–2), 46–53.
- Bogin, B., Silva, M.I.V., Rios, L., 2007. Life history trade-offs in human growth: adaptation or pathology? *Am. J. Hum. Biol.* 19 (5), 631–642.
- Brooks, S., Suchey, J.M., 1990. Skeletal age determination based on the *os pubis*: a comparison of the acsádi-nemeskéri and suchey-brooks methods. *Hum. Evol.* 5 (3), 227–238.
- Brown, T.A., Nelson, D.E., Vogel, J.S., Southon, J.R., 1988. Improved collagen extraction by modified longin method. *Radiocarbon* 30 (2), 171–177.
- Buikstra, J., Ubelaker, D.H., 1994. Standards for data collection from human skeletal remains. *Arkansas Archaeological Survey, Fayetteville, Arkansas*.
- Bush, H.M., 1989. The Recognition of Physiological Stress in Human Skeletal Material: A Critique of Method and Theory with Specific Reference to the Vertebral Column. Ph.D. Dissertation. University of Sheffield.
- Calder, P.C., 2013. Feeding the immune system. *Proc. Nutr. Soc.* 72 (3), 299–309.
- Calder, P.C., Jackson, A.A., 2000. Malnutrition, infection and immune function. *Nutr. Res. Rev.* 13 (1), 3–29.
- Conde, M.A.S., 1996. Tomar Medieval. O espaço e os homens, Cascais.
- Curto, A., Maurer, A.F., Barrocas-Dias, C., Mahoney, P., Fernandes, T., Fahy, G.E., 2018. Did military orders influence the general population diet? Stable isotope analysis from Medieval Tomar, Portugal. *Archaeological and Anthropological Sciences* 1–13.
- D'Ortenzio, L., Brickley, M., Schwarcz, H., Prowse, T., 2015. You are not what you eat during physiological stress: isotopic evaluation of human hair. *Am. J. Phys. Anthropol.* 157 (3), 374–388.
- Deitch, E.A., 1990. The role of intestinal barrier failure and bacterial translocation in the development of systemic infection and multiple organ failure. *Arch. Surg.* 125 (3), 403–404.
- DeNiro, M.J., Epstein, S., 1976. You are what you eat (plus a few‰): the carbon isotope cycle in food chains. *Geological Society of America* 6, 834.
- DeNiro, M.J., Epstein, S., 1978. Influence of diet on the distribution of carbon isotopes in animals. *Geochem. Cosmochim. Acta* 42 (5), 495–506.
- Deschner, T., Fuller, B.T., Oelze, V.M., Boesch, C., Hublin, J., Mundry, R., Hohmann, G., 2012. Identification of energy consumption and nutritional stress by isotopic and elemental analysis of urine in bonobos (*Pan paniscus*). *Rapid Commun. Mass Spectrom.* 26 (1), 69–77.
- DeWitte, S.N., 2010. Sex differentials in frailty in medieval England. *Am. J. Phys. Anthropol.* 143 (2), 285–297.
- DeWitte, S.N., Hughes-Morey, G., 2012. Stature and frailty during the Black Death: the effect of stature on risks of epidemic mortality in London, AD 1348–1350. *J. Archaeol. Sci.* 39 (5), 1412–1419.
- DeWitte, S.N., Stojanowski, C.M., 2015. The osteological paradox 20 years later: past perspectives, future directions. *J. Archaeol. Res.* 23 (4), 397–450.
- DeWitte, S.N., Wood, J.W., 2008. Selectivity of Black Death mortality with respect to pre-existing health. *Proc. Natl. Acad. Sci. Unit. States Am.* 105 (5), 1436–1441.
- Fuller, B.T., Fuller, J.L., Sage, N.E., Harris, D.A., O'Connell, T.C., Hedges, R.E., 2005. Nitrogen balance and  $\delta^{15}\text{N}$ : why you're not what you eat during nutritional stress. *Rapid Commun. Mass Spectrom.* 19 (18), 2497–2506.
- Gaye-Siesseger, J., Focken, U., Abel, H., Becker, K., 2004. Individual protein balance strongly influences  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  values in Nile tilapia, *Oreochromis niloticus*. *Naturwissenschaften* 91 (2), 90–93.
- Gonçalves, I., 2004. Entre a abundância e a miséria: as práticas alimentares da Idade Média portuguesa. *Estudos medievais. Quotidiano Medieval: imaginário, representação e práticas*. Livros Horizonte, Lisboa.
- Goodman, A.H., Martin, D.L., 2002. Reconstructing Health Profiles from Skeletal Remains. *The Backbone of History*. Cambridge University Press, Cambridge, UK.
- Grossman, C.M., Sappington, T.S., Burrows, B.A., Lavietes, P.H., Peters, J.P., 1945. Nitrogen metabolism in acute infections. *J. Clin. Invest.* 24 (4), 523–531.
- Heaton, T.H.E., 1987. The  $^{15}\text{N}/^{14}\text{N}$  ratios of plants in South Africa and Namibia: relationship to climate and coastal/saline environments. *Oecologia* 74, 236–246.
- Heaton, T.H.E., Vogel, J.C., von la Chevallerie, G., Collett, G., 1986. Climatic influence on the isotopic composition of bone nitrogen. *Nature* 322, 822–823.
- Hedges, R.E., Clement, J.G., Thomas, C.D.L., O'Connell, T.C., 2007. Collagen turnover in the adult femoral mid-shaft: modeled from anthropogenic radiocarbon tracer measurements. *Am. J. Phys. Anthropol.* 133 (2), 808–816.
- Hobson, K.A., Alisauskas, R.T., Clark, R.G., 1993. Stable-nitrogen isotope enrichment in avian tissues due to fasting and nutritional stress: implications for isotopic analyses of diet. *Condor* 388–394.
- Huss-Ashmore, R., Schall, J., Hediger, M., 1992. Health and Lifestyle Change. UPenn Museum of Archaeology.
- Katzenberg, M.A., Lovell, N.C., 1999. Stable isotope variation in pathological bone. *Int. J. Osteoarchaeol.* 9 (5), 316–324.
- Keusch, G.T., Farthing, M.J., 1986. Nutrition and infection. *Annu. Rev. Nutr.* 6 (1), 131–154.
- Kiple, K.F., 1993. The treponematoses. In: Kiple, K.F. (Ed.), *The Cambridge World History of Human Disease*. Cambridge University Press, Cambridge, pp. 1053–1055.
- Kjellström, A., Storå, J., Possnert, G., Linderholm, A., 2009. Dietary patterns and social structures in Medieval Sigtuna, Sweden, as reflected in stable isotope values in human skeletal remains. *J. Archaeol. Sci.* 36 (12), 2689–2699.
- Klaus, H.D., 2014. Frontiers in the bioarchaeology of stress and disease: cross-disciplinary perspectives from pathophysiology, human biology, and epidemiology. *Am. J. Phys. Anthropol.* 155 (2), 294–308.
- Kuvibidila, S., Yu, L., Ode, D., Warrier, R.P., 1993. The immune response in protein-energy malnutrition and single nutrient deficiencies. In: *Nutrition and Immunology*. vols. 121–155 Springer US.
- Larsen, C.S., 2002. Bioarchaeology: the lives and lifestyles of past people. *J. Archaeol. Res.* 10 (2), 119–166.
- Lee, W.H., Woodward, B.D., 1996. The CD4/CD8 ratio in the blood does not reflect the response of this index in secondary lymphoid organs of weaning mice e in models of protein-energy malnutrition known to depress thymus-dependent immunity. *J. Nutr.* 126 (4), 849–859.
- Linderholm, A., Kjellström, A., 2011. Stable isotope analysis of a medieval skeletal sample indicative of systemic disease from Sigtuna Sweden. *J. Archaeol. Sci.* 38 (4), 925–933.
- Linderholm, A., Jonson, C.H., Svensk, O., Liden, K., 2008. Diet and status in Birka: stable isotopes and grave goods compared. *Antiquity* 82, 446–461.
- Longin, R., 1971. New method of collagen extraction for radiocarbon dating. *Nature* 230 (5291), 241–242.
- Lovejoy, C.O., Meindl, R.S., Pryzbeck, T.R., Mensforth, R.P., 1985. Chronological metamorphosis of the auricular surface of the ilium: a new method for the determination of adult skeletal age at death. *Am. J. Phys. Anthropol.* 68 (1), 15–28.
- Martorell, R., Yarbrough, C., Yarbrough, S., Klein, R.E., 1980. The impact of ordinary illnesses on the dietary intakes of malnourished children. *Am. J. Clin. Nutr.* 33 (2), 345–350.
- Mata, L.J., Urrutia, J.J., Lechtig, A., 1971. Infection and nutrition of children of a low socioeconomic rural community. *Am. J. Clin. Nutr.* 24 (2), 249–259.
- Minagawa, M., Wada, E., 1984. Stepwise enrichment of  $^{15}\text{N}$  along food chains: further evidence and the relation between  $\delta^{15}\text{N}$  and animal age. *Geochem. Cosmochim. Acta* 48 (5), 1135–1140.

- Mitra, A.K., Akramuzzaman, S.M., Fuchs, G.J., Rahman, M.M., Mahalanabis, D., 1997. Long-term oral supplementation with iron is not harmful for young children in a poor community of Bangladesh. *J. Nutr.* 127 (8), 1451–1455.
- Müldner, G., Richards, M.P., 2007. Stable isotope evidence for 1500 years of human diet at the city of York, UK. *Am. J. Phys. Anthropol.* 133 (1), 682–697.
- Müldner, G., Montgomery, J., Cook, G., Ellam, R., Gledhill, A., Lowe, C., 2009. Isotopes and individuals: diet and mobility among the medieval Bishops of Whithorn. *Antiquity* 83 (322), 1119–1133.
- Oelbermann, K., Scheu, S., 2001. Stable isotope enrichment ( $\delta^{15}\text{N}$  and the  $\delta^{13}\text{C}$ ) in a generalist predator (*Pardosalugubris*, Araneae: lycosidae): effects of prey quality. *Oecologia* 130, 337–344.
- Ortner, D.J., 2003. *Identification of Pathological Conditions in Human Skeletal Remains*. Academic Press.
- Ortner, D.J., Putschar, W.G.J., 1985. *Identification of Pathological Conditions in Human Skeletal Remains*. Smithsonian Institution Press, Washington, London.
- Paine, R.R., Brenton, B.P., 2006. The paleopathology of pellagra: investigating the impact of prehistoric and historical dietary transitions to maize. *Journal of Anthropological Sciences* 84, 125–135.
- Peck, J.J., 2013. Status, health, and lifestyle in middle iron age Britain: a bioarchaeological study of elites and non-elites from East Yorkshire, Northern England. *International Journal of Paleopathology* 3 (2), 83–94.
- Phenice, T.W., 1969. A newly developed visual method of sexing the os pubis. *Am. J. Phys. Anthropol.* 30 (2), 297–301.
- Pinhasi, R., Mays, S. (Eds.), 2008. *Advances in Human Palaeopathology*. John Wiley & Sons.
- Polet, C., Katzenberg, M.A., 2003. Reconstruction of the diet in a mediaeval monastic community from the coast of Belgium. *J. Archaeol. Sci.* 30 (5), 525–533.
- Powanda, M.C., 1977. Changes in body balances of nitrogen and other key nutrients: description and underlying mechanisms. *Am. J. Clin. Nutr.* 30 (8), 1254–1268.
- Ragsdale, B.D., Lehmer, L.M., 2012. A knowledge of bone at the cellular (histological) level is essential to paleopathology. In: Grauer, A.L. (Ed.), *A Companion to Paleopathology*. Wiley-Blackwell, Chichester, UK, pp. 227–249.
- Redlich, K., Smolen, J.S., 2012. Inflammatory bone loss: pathogenesis and therapeutic intervention. *Nat. Rev. Drug Discov.* 11, 234–250.
- Reitsema, L.J., Vercellotti, G., 2012. Stable isotope evidence for sex-and status-based variations in diet and life history at medieval Trino Vercellese, Italy. *Am. J. Phys. Anthropol.* 148 (4), 589–600.
- Reitsema, L.J., Crews, D.E., Polcyn, M., 2010. Preliminary evidence for medieval Polish diet from carbon and nitrogen stable isotopes. *J. Archaeol. Sci.* 37 (7), 1413–1423.
- Reitsema, L.J., Vercellotti, G., Boano, R., 2016. Subadult dietary variation at Trino Vercellese, Italy, and its relationship to adult diet and mortality. *Am. J. Phys. Anthropol.* 160 (4), 653–664.
- Richards, M.P., Hedges, R.E., 1999. Stable isotope evidence for similarities in the types of marine foods used by Late Mesolithic humans at sites along the Atlantic coast of Europe. *J. Archaeol. Sci.* 26 (6), 717–722.
- Robb, J., Bigazzi, R., Lazzarini, L., Scarsini, C., Sonogo, F., 2001. Social “status” and biological “status”: a comparison of grave goods and skeletal indicators from pontecagnano. *Am. J. Phys. Anthropol.* 115 (3), 213–222.
- Salamon, M., Coppa, A., McCormick, M., Rubini, M., Vargiu, R., Tuross, N., 2008. The consilience of historical and isotopic approaches in reconstructing the medieval Mediterranean diet. *J. Archaeol. Sci.* 35 (6), 1667–1672.
- Scheuer, J.L., Black, S., 2000. Development and ageing of the juvenile skeleton. In: Cox, M., Mays, S. (Eds.), *Human Osteology in Archaeology and Forensic Science*. vols. 9–21 Greenwich Medical Media, London.
- Schoeninger, M.J., DeNiro, M.J., 1984. Nitrogen and carbon isotopic composition of bone collagen from marine and terrestrial animals. *Geochem. Cosmochim. Acta* 48 (4), 625–639.
- Schoeninger, M.J., DeNiro, M.J., Tauber, H., 1983. Stable nitrogen isotope ratios of bone collagen reflect marine and terrestrial components of prehistoric human diet. *Science* 220 (4604), 1381–1383.
- Schutkowski, H., Herrmann, B., Wiedemann, F., Bocherens, H., Grupe, G., 1999. Diet, status and decomposition at Weingarten: trace element and isotope analyses on early mediaeval skeletal material. *J. Archaeol. Sci.* 26 (6), 675–685.
- Scorrano, G., Brillì, M., Martínez-Labarga, C., Giustini, F., Pacciani, E., Chilleri, F., Scaldaferrì, F., Gasbarrini, A., Gasbarrini, G., Rickards, O., 2014. Palaeodiet reconstruction in a woman with probable celiac disease: a stable isotope analysis of bone remains from the archaeological site of Cosa (Italy). *Am. J. Phys. Anthropol.* 154 (3), 349–356.
- Scrimshaw, N.S., SanGiovanni, J.P., 1997. Synergism of nutrition, infection, and immunity: an overview. *Am. J. Clin. Nutr.* 66 (2), 464S–477S.
- Sealy, J.C., van der Merwe, N.J., Thorp, J.A., Lanham, J.L., 1987. Nitrogen isotopic ecology in southern Africa: implications for environmental and dietary tracing. *Geochem. Cosmochim. Acta* 51, 2707–2717.
- Sherman, P., Forstner, J., Roomi, N., Khatri, I., Forstner, G., 1985. Mucin depletion in the intestine of malnourished rats. *Am. J. Physiol. Gastrointest. Liver Physiol.* 248 (4), G418–G423.
- Siek, T., 2013. The osteological paradox and issues of interpretation in paleopathology. *Vis-à-Vis: Explor. Anthropol.* 13, 92–101.
- Sponheimer, M., Robinson, T., Ayliffe, L., Roeder, B., Hammer, J., Passey, B., et al., 2003. Nitrogen isotopes in mammalian herbivores: hair  $\delta^{15}\text{N}$  values from a controlled feeding study. *Int. J. Osteoarchaeol.* 13 (1–2), 80–87.
- Steckel, R.H., Larsen, C.S., Sciuilli, P.W., Walker, P.L., 2006. Data collection codebook. The Global History of Health Project 1–41.
- Steele, K., Daniel, R.M., 1978. Fractionation of nitrogen isotopes by animals: a further complication to the use of variations in the natural abundance of  $^{15}\text{N}$  for tracer studies. *J. Agric. Sci.* 90 (1), 7–9.
- Temple, D.H., Goodman, A.H., 2014. Bioarcheology has a “health” problem: conceptualizing “stress” and “health” in bioarcheological research. *Am. J. Phys. Anthropol.* 155 (2), 186–191.
- Tieszen, L.L., Boutton, T.W., Tesdahl, K.G., Slade, N.A., 1983. Fractionation and turnover of stable carbon isotopes in animal tissues: implications for  $\delta^{13}\text{C}$  analysis of diet. *Oecologia* 57 (1–2), 32–37.
- Turner-Walker, G., 2008. The chemical and microbial degradation of bones and teeth. *Advances in human palaeopathology* 592.
- Ulijaszek, S.J., Mann, N., Elton, S., 2012. *Evolving Human Nutrition: Implications For Public Health*. Cambridge University Press.
- Valente, J., 1998. The new frontier: the role of the knights templar in the establishment of Portugal as an independent kingdom. *Mediterr. Stud.* 7, 49–65.
- Van der Merwe, N.J., Vogel, J.C., 1978.  $^{13}\text{C}$  content of human collagen as a measure of prehistoric diet in woodland North America. *Nature* 276 (5690), 815.
- Vicente, M., 2013. *Entre Zêzere e Tejo: Propriedade e Povoamento*. Doutoramento em História Medieval. (Universidade de Lisboa [Unpublished]).
- Watts, R., 2013. Childhood development and adult longevity in an archaeological population from Barton-upon-Humber, Lincolnshire, England. *International Journal of Paleopathology* 3 (2), 95–104.
- Welle, S., 1999. *Human Protein Metabolism*. Springer, Berlin Heidelberg New York.
- Weston, D.A., 2008. Investigating the specificity of periosteal reactions in pathology museum specimens. *Am. J. Phys. Anthropol.* 137 (1), 48–59.
- Weston, D.A., 2009. Brief communication: paleohistopathological analysis of pathology museum specimens: can periosteal reaction microstructure explain lesion etiology? *Am. J. Phys. Anthropol.* 140 (1), 186–193.
- Weston, D.A., 2012. Nonspecific infection in paleopathology: interpreting periosteal reactions. In: Grauer, A.L. (Ed.), *A Companion to Paleopathology*. Wiley-Blackwell, Chichester, UK, pp. 492–512.
- Wood, J., Milner, G., Harpending, H., Weiss, K., 1992. The osteological paradox - problems of inferring prehistoric health from skeletal samples. *Curr. Anthropol.* 33 (4), 343–370.
- Woodward, B., 1998. Protein, calories, and immune defences. *Nutr. Rev.* 56 (1), S84–S92.
- Woodward, B., 2001. The effect of protein-energy malnutrition on immune competence. In: Suskind, R.M., Tontisirin, K.T. (Eds.), *Nestlé Nutrition Workshop Series*. vol. 45. Lippincott-Raven, Philadelphia, pp. 89–120.
- Woodward, B.D., Miller, R.G., 1991. Depression of thymus-dependent immunity in wasting protein-energy malnutrition does not depend on an altered ratio of helper (CD4+) to suppressor (CD8+) T cells or on a disproportionately large atrophy of the T-cell relative to the B-cell pool. *Am. J. Clin. Nutr.* 53 (5), 1329–1335.