



Rift-related volcanism predating the birth of the Rheic Ocean (Ossa-Morena zone, SW Iberia)

T. Sánchez-García ^{a,*}, F. Bellido ^a, M.F. Pereira ^b, M. Chichorro ^c, C. Quesada ^a, Ch. Pin ^d, J.B. Silva ^e

^a IGME, c/Ríos Rosas, 23, 28003-Madrid, Spain

^b Dpto. Geociências, Centro de Geofísica de Évora, Univ. Évora, Apt. 94, 7001-554 Évora, Portugal

^c CICEGe - Centro de Investigação em Ciência e Engenharia Geológica, Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, Quinta da Torre, 2829-516 Caparica, Portugal

^d Dpt. Géologie, CNRS, Univ. Blaise Pascal, rue Kessler, 63038 Clermont-Ferrand, France

^e Dpto. Geologia, Faculdade Ciências, Univ. Lisboa, Portugal

ARTICLE INFO

Article history:

Received 6 May 2009

Received in revised form 26 September 2009

Accepted 13 October 2009

Available online 25 October 2009

Keywords:

Rift magmatism
Cambrian–Ordovician
Geochemistry
Rheic Ocean
Ossa-Morena zone
Sm–Nd isotopes

ABSTRACT

Two very different periods of magma emplacement in the crust of the Ossa-Morena zone (early and main events) in SW Iberia have been previously interpreted to record a Cambrian/Early Ordovician rifting event that is thought to have culminated in the opening of the Rheic Ocean during the Early Ordovician. New stratigraphic, petrographic, geochemical and Sm–Nd isotope data from Cambrian volcanic rocks included in six key low-grade sections in both Portugal and Spain considerably improve our understanding of these events. These data: (1) confirm the existence of two rift-related magmatic events in the Cambrian of the Ossa-Morena zone, (2) demonstrate that the early rift-related event was associated with migmatite and core-complex formation in the mid-upper crust and is represented by felsic peraluminous rocks, the parent magmas of which were derived mainly from crustal sources, and (3) show the main rift-related event to be represented by a bimodal association of felsic and mafic rocks with minor amounts of intermediate rocks. Some of the mafic rocks show N-MORB affinity, whereas others have OIB or E-MORB affinities, suggesting different heterogeneous mantle sources (depleted and enriched, asthenospheric and lithospheric, plume-like and non-plume-like). The acid and intermediate rocks appear to represent hybrid mixtures of crust and mantle-derived magmas.

This new data supports the hypothesis that the onset of rifting was associated with a process of oblique ridge-trench collision. We interpret the significant differences between the early and main events as reflecting the evolution from a wide rift stage with passive extension mainly accommodated by lower-crust flow in a high heat-flow setting, to a narrow rift stage with active extension characterized by extension rates that outpaced thermal diffusion rates.

© 2009 International Association for Gondwana Research. Published by Elsevier B.V. All rights reserved.

1. Introduction

Two events played crucial roles in the evolution of the Iberian Massif during the Paleozoic. The first of these was a Cambrian/Early Ordovician rifting event, which was largely responsible for the compartmentalization of the Paleozoic Iberian autochthonous margin of Gondwana and consequent paleogeographic and lithotectonic differences between the Cantabrian, West Asturian-Leonese, Central Iberian, and Ossa-Morena zones (Fig. 1) (Liñán and Quesada, 1990; Quesada et al., 1991; Quesada, 1991; Sánchez-García, 2001; Sánchez-García et al., 2003; Quesada, 2006; Quesada et al., 2006; Sánchez-García et al., 2008a). This rifting event is interpreted as having culminated in the sequential opening of the Rheic Ocean by the Early Ordovician (Quesada, 1991; Sánchez-García et al., 2003, 2008a; Fernández et al., 2008). The second event was the Variscan orogeny, which was caused by continental collision following the closure of the Rheic Ocean and is responsible for the

deformation of the Paleozoic Iberian margin of Gondwana and the present geometrical arrangement of units (Fig. 1) (Ribeiro et al., 1990; Quesada et al., 1991; Quesada, 1991; Quesada et al., 2006; Quesada, 2006).

Previous to these Paleozoic events, the geodynamic evolution of northern Gondwana was characterized by a period of arc growth and accretion to the continental margin during the Ediacaran (Cadomian orogeny). This process had come to a fairly abrupt halt by the Early Cambrian, when the margin, including parts of the previously accreted Cadomian-Avalonian arc (Quesada, 1991; Nance et al., 2002; Sánchez-García et al., 2003; Murphy et al., 2006; Pereira and Quesada, 2006; Pereira et al., 2006; Sánchez-García et al., 2008a), started to undergo differential uplift and erosion together with a cessation of subduction-related magmatism. This was followed by an onset of extensional deformation, which was responsible for crustal thinning with compartmentalization into subsiding (graben) and more stable (horst) domains (Liñán and Quesada, 1990). At the same time mantle upwelling triggered rift-related igneous activity, which reached its maximum expression in Iberia within the Ossa-Morena zone (Sánchez-García et al., 2003, 2008a). This evolution from subduction/arc growth to continental

* Corresponding author.

E-mail address: t.sanchez@igme.es (T. Sánchez-García).