

The management and preservation of communitarian interest habitats in the Natural Park of Serra da Estrela (Portugal)

António Meireles⁽¹⁾, Rafael Neiva⁽²⁾, Isabel Passos⁽¹⁾, Carlos Vila-Viçosa⁽¹⁾, Rodrigo Pinava-Ferreira⁽¹⁾ & Carlos Pinto-Gomes⁽¹⁾

⁽¹⁾ Parque Natural da Serra da Estrela, Rua 1º de Maio, nº 2, P-6260-101 Manteigas; ⁽²⁾ meireles@portugalmail.pt

⁽¹⁾ Universidade de Évora, Departamento de Ecologia, Rua Romão Ramalho, nº 59, P-7000-671 Évora

Abstract. - In this paper we present the management and preservation strategy of communitarian interest habitats which can be found in the Natural Park of Serra da Estrela (PNSE), Portugal. After presenting a general characterization of PNSE, where the main biophysical aspects of the territory are signalled, the description of the main work carried out in this Protected Area, in terms of characterization, improvement, diffusion and management of habitats of communitarian interest to be found here, will follow.

Key words: - Serra da Estrela - habitats - Natura 2000 - management and preservation.

Résumé. - On présente ici la stratégie de gestion et de conservation des habitats d'intérêt communautaire du Parc national de la Sierra de Estrela (PNSE), Portugal. Après une présentation générale biophysique de ce parc, vient la description des principaux travaux menés sur cette aire protégée en termes de caractérisation, amélioration, diffusion et gestion de ces habitats.

Mots clés: - Serra da Estrela - habitats - Natura 2000 - gestion et conservation.

I. INTRODUCTION

Being in mind the goal of favouring the maintenance of biodiversity of its member states, in a sustainable development perspective, the European Union, started, in the late eighties, a process that would culminate in the creation of an european ecological net, commonly known as Natura 2000.

The selection of places to integrate the net of natural spaces, was made through two communitarian directives, known as Birds Directive and Habitats Directive, whose transposition into Portuguese legislation was made on 24th April 1999, through a single document, the law by decree (Decreto-Lei) nº 140/99, which in its turn was amended on 24th February 2005, by the law by decree (Decreto-Lei) nº 49/2005.

Wolock C.E., M. Jebb & C.B. Osmond, 1994.- Photoinhibition and recovery in tropical plant species: response to disturbance. *Oecologia*, **97**, 297-307.

APA, 1994.- *Métodos oficiais de análise de solos y aguas*. Ministerio de Agricultura, Pesca y Alimentación (España), Madrid, 182 p.

Artinez-González M.A., A. Sánchez-Villegas & J. Faulin (eds.), 2006.- *Bioestadística amigable*. Ed. Diaz de Santos, Spain, 919 p.

Arnwell K. & G.N. Johnson, 2000.- Chlorophyll fluorescence, a practical guide. *J. Exp. Bot.*, **51**, 659-668.

Finander O., S. Somersalo, T. Holopainen & R.J. Strasser, 1996.- Scots pine after exposure to elevated ozone and carbon dioxide probed by reflectance spectra and chlorophyll a fluorescent transients. *J. Plant Physiol.*, **148**, 229-236.

Flo M.E., M.L. Rodríguez-Tamayo, M.L. Jiménez, J.F. Mota, 2001.- Recapitulación sobre el comportamiento bioquímico de algunos gipsófitos y halófitos ibéricos. *Monogr. Flora y Vegetación Béticas*, **12**, 77-95.

Plina A., J. Loidi & F. Fernández-González, 1993.- Sobre las comunidades de matorral de la depresión del Ebro (España). *Bot. Complut.*, **18**, 11-50.

Ja J.F., 2001.- Análisis de agrupamientos o de "cluster" aplicado al estudio de la flora y vegetación de yesos. In: *Aplicaciones ambientales del análisis multivariante*. P. Aguilera & A. Garrido (eds.), Universidad de Almería, 143-176.

Ja J.F., J.J. Alvarado, F. Gómez, F. Valle & J. Cabello, 1995.- Vegetación gipsícola y conservación de la naturaleza. *Coll. Phytosoc.*, **21**, 677-688.

Ja J.F., J. Cabello, M. Cueto, F. Gómez, E. Giménez & J. Peñas, 1997.- Datos sobre la vegetación del sur-este de Almería. *Desierto de Tabernas, Karst en Yesos de Sorbas y Cabo de Gata*. Universidad de Almería, 130 p.

Ja J.F., M.E. Merlo, F. Pérez-García, J.A. Garrido, F. Martínez-Hernández, P. Sánchez-Gómez, M.L. Rodríguez-Tamayo, L. Posadas, S. Ruiz & C. Gil de Carrasco, 2007.- Plants growing on unusual substrates: the edaphism. Notes from the antipodes (Iberian Peninsula). 49th IAVS Conference Abstracts Book, New Zealand, 165.

Ja J.F., A. Sola, E.D. Diana & M.L. Jiménez-Sánchez, 2003.- Plant succession in abandoned gypsum quarries in SE Spain. *Phytocoenologia*, **33** (1), 13-28.

Ja J.F., A.J. Sola, M.L. Jiménez-Sánchez, F.J. Pérez-García & M.E. Merlo, 2004a.- Gypsicolous flora, conservation and restoration of quarries in the southeast of the Iberian Peninsula. *Biodiv. Conserv.*, **13**, 1797-1808.

Ja J.F., A.J. Sola, M.L. Jiménez-Sánchez, F.J. Pérez-García & M.E. Merlo, 2004b.- Gypsophilous flora and quarries: some remarks on the preservation and restoration of gypsum outcrops. *Ecologia Medit.*, **30** (2), 107-109.

Ilkey S.S. & R.W. Pearcy, 1992.- Interaction between acclimation and photoinhibition of photosynthesis of a tropical understory herb, *Alocasia macrorrhiza*, during simulated canopy gap formation. *Funct. Ecol.*, **6**, 719-729.

Ochoa M.J., 1982.- Relaciones entre el medio y comunidades vegetales del sabinar continental árido del valle del Ebro. *Com. INIA*, ser. Rec. Nat., **14**, 1-52.

Oliveira G. & J. Peñuelas, 2000.- Comparative photochemical and phenomorphological responses to winter stress of an evergreen (*Quercus ilex* L.) and a semi-deciduous (*Cistus albidus* L.) Mediterranean woody species. *Acta Oecol.*, **21**, 97-107.

Palik B.J., K.W. Michener, R.J. Mitchell & J.W. Jones, 1999.- The effect of landform and plant size on mortality and recovery of longleaf pine during a 100-year flood. *Ecoscience*, **6** (2), 255-263.

Parsons R.F., 1976.- Gypsophily in plants, a review. *Am. Midl. Nat.*, **96** (1), 1-20.

Pérez-Pujalte A. & C. Oyonarte, 1987.- Mapa de suelos 1: 100.000. Tabernas, Hoja 1030, Proyecto LUCDEME, ICONA, 115 p.

Pérez-Pujalte A. & C. Oyonarte, 1988.- Mapa de suelos 1: 100.000. Sorbas, Hoja 1031, Proyecto LUCDEME, ICONA, 148 p.

Porta J., 1998.- Methodologies for analysis and characterization of gypsum in soils: a review. *Geoderma*, **87**, 31-46.

Richard L.A. & L.R. Weaver, 1969.- Soil water. In: *Diagnosis and improvement of saline and alkali soils*. L.R. Richard (ed.), *Agriculture Handbook*, **60**, United States Dept. Agriculture, WashingtonDC.

Rivas-Martínez S., T.E. Díaz, F. Fernández-González, J. Izco, J. Loidi, M. Lousá & A. Penas, 2002.- Vascular plant communities of Spain and Portugal. *Itinerario Geobot.*, **15** (1), 1-432.

Schreiber U. & W. Bilger, 1993.- Progress in chlorophyll fluorescence research: major developments during the past years in retrospect. *Progress Bot.*, **54**, 151-173.

Schreiber U., H. Hormann, C. Neubauer, & C. Klughammer, 1995.- Assessment of photosystem II photochemical quantum yield by chlorophyll fluorescence quenching analysis. *Austr. J. Plant Physiol.*, **22**, 209-220.

Sokal R.R. & F.J. Rohlf, 1995.- *Biometry*, 3rd. W.H. Freeman and Company, NY.

Sola A.J., M.L. Rodríguez-Tamayo, F.J. Pérez-García, E. Dana, M. Cerrillo & J.F. Mota, 2001.- Caracterización florística y ecológica de una catena de vegetación en los aljzares de Sorbas (Almería). In: *Valoración y Gestión de Espacios Naturales*. E. Cano, A. García Fuentes, J.A. Torres & C. Salazar (eds.), Servicio de Publicaciones de la Universidad de Jaén, 349-354.

Tyurin I.V., 1951.- Analytical procedure for comparative study of soil humus. *Trudy Pochv. Inst. Dokuchaev*, **38**, 5.

Acknowledging the importance of natural and semi-natural habitats to the preservation of European biodiversity, these have taken on an essential role in the process of selection of the places to integrate the future Natura 2000. Therefore, one of the selection criteria for these areas was the existence, at the site, of habitats which are contained in the Annex II of the Directive Habitats.

Due to its richness and biological diversity, made up of endemic species and very rare habitats nation wide, a large part of the massif of Serra da Estrela was included in the National Sites List and put forward for consideration to integrate the future Natura 2000 (Site PTCON0016).

Despite the fact that many botanists went through Serra da Estrela and there is work done on habitats, previously carried out in the scope of the Natura 2000 (Jansen, 1997), there was, until 2006, no habitat mapping of habitats that could allow, among other things, the integrated preservation and management of this biological heritage which exists in the ranked area. In this sense, one of the most recent strategies of the PNSE, consisted in identifying and establishing boundaries for communitarian interest habitats present in its scope, done in cooperation with the Instituto Superior Agrário de Bragança (Agrarian Superior Institute of Bragança). Next, establishing a partnership with the Universidade de Évora (Évora University), a work of improvement, diffusion and proposals of management measures for the plant community heritage found here was carried out.

II. CHARACTERISATION OF THE NATURAL PARK OF SERRA DA ESTRELA

Located in Centre-East of Portugal, in the extension of the Central System through Portuguese territory, Serra da Estrela is a mountainous massif of national reference, for it includes the highest point of continental Portugal. Thus, many studies have been carried out in this territory (e.g. Henriques, 1883; Sampaio, 1910; Braun-Blanquet *et al.*, 1952, 1956, 1964; Malato-Beliz, 1955; Du vigneaud, 1962; Rivas-Martínez, 1974; Pinto da Silva *et al.*, 1976; Pinto da Silva & Teles, 1986; Jansen, 2002), having had their national importance been recognised in 1976 through the creation of the Natural Park of Serra da Estrela (PNSE), by the law by decree (Decreto-Lei) n° 557/76, dated of July 16th. At this time, the protection and the preservation of this mountainous system were defined as prime objectives, not only as far as natural values are concerned, but also for the maintenance and diffusion of traditional practices and local culture.

The PNSE is a vast protected area that extends for about one hundred thousand hectares, in the districts of Castelo Branco and Guarda. From the biogeographical point of view it is included, mainly, in the territories of the Estrelense sector, of the Carpetan-Leonese (Rivas-Martínez, 1987; Costa *et al.*, 1998; Rivas-Martínez *et al.*, 2002). Besides the most diffused natural values, as the snow and the very beautiful landscapes, Serra da Estrela also includes an enormous biological heritage.

The current diversity of fauna and flora found here is also due to several factors such as the geographical position, the topographical complexity, the geological history of the territory, the climatic peculiarities and the secular action of man. Among these, the great variety of physical environments, mostly when it comes to climate, takes on a particularly important role. Even though a great part of the area is under a climate of Mediterranean type, in the areas of greatest altitude and in the slopes more to the west, we can already see the transition to a climate of moderate influence. Besides this variation in latitude, there are also altitude related variations of the climate, by which it is possible to observe three

distinct bioclimatic levels. In the less high areas (until 850 metres) there is a basal level, hotter and drier, with bioclimatics belts of mesomediterranean or mesotemperate type. In this level the climatic vegetation corresponds to mixed woods of evergreen and deciduous *Quercus*, dominated by *Quercus suber* (in the area of Mediterranean influence), or deciduous woods of *Quercus robur* (in the area of atlantic influence). Above this level, between the (850) 900 metres the climate becomes the Supramediterranean or Supratemperate and the climatic vegetation corresponds to woods dominated by *Quercus pyrenaica*. Finally, in the upper level, located above 1600 (1800 metres) and climatologically more severe (Oromediterranean or Submediterranean orotemperate), we find the climatic domination of the juniper of *Juniperus communis* subsp. *nana*.

In geological terms, Serra da Estrela is a massif of recent formation, mostly granitic, holding evidence of the last great glaciations.

III. THE HABITATS OF COMMUNITARIAN INTEREST TO BE FOUND AT NATURAL PARK OF SERRA DA ESTRELA

Currently, there are 32 habitats of communitarian interest referenced for the PNSE, five of which are a priority as to preservation (Table I). Of these, mainly the ones found in the upper plateau, some of them of rarity or even exclusive to Portugal, stand out.

IV. CHARACTERISATION, IMPROVEMENT, DIFFUSION AND MANAGEMENT OF HABITATS IN NATURAL PARK OF SERRA DA ESTRELA

Having as the final goal the preservation and improvement of the habitats of communitarian interest in Serra da Estrela, in the last years, several studies, which allowed to know, map, improve and begin a work of management of this biological heritage, were carried out.

This process can be divided into four different phases. In a first phase the habitats found in the territory were identified and mapped. This process was carried out through cooperation between the PNSE and the Agrarian Superior Institute of Bragança, which led to the attainment, in the end, of a mapping of the natural and semi-natural habitats, of communitarian interest found in the PNSE (Esteves *et al.*, 2006). Besides this, a work of mapping the habitats of one of the most emblematic areas in the Park, the Glacier Valley of the Zêzere River, had previously been done. This work, carried out by Passos (2005), of the University of Évora, gave the knowledge, in detail, of the habitats found in this area.

Based in these mappings, as well as in the mapping of soil use, also made at the same time in a cooperation between the Agrarian Superior Institute of Bragança and the PNSE, the next step was taken: the territorial improvement of the PNSE, from the point of view of the plant community (Meireles *et al.*, 2006a). Such work had as prime objective the finding of territories and vegetable formations priority to preservation, in a perspective of territory management. The used methodology was that put forward by Maeza & Cadiñanos (2000). It gave the possibility of getting to a numeric value capable of translating the interest of preserving some of the main vegetable communities found in PNSE. The results show the « exceptional » value of some of the woods of the region, with a very rare and separate distribution presently. In this position we find the climatic oak-groves of *Quercus pyrenaica* (habitat 9230), the woods of *Prunus lusitanica* (habitat 5230*) and the

Table 1.- List of habitats of communitarian interest to be found in the Natural Park of Serra da Estrela.
 Tableau 1.- Liste des habitats d'intérêt communautaire du Parc national de la Sierra de Estrela.

Code	Name	Code	Name
3130	Oligotrophic to mesotrophic standing waters with vegetation of the <i>Littorelletta uniflorae</i> and/or <i>Isostio-Nanojuncetea</i>	6410	Molinia meadows on calcareous, peaty or clayey -siltadensols (<i>Molinia caerulea</i>)
3150	Natural eutrophic lakes with <i>Megnopotamion</i> or <i>Hydrocharition</i> - type vegetation	6430	Hydrophilous tall herb fringe communities of plains and of the montane to alpine levels
3260	Water courses of plain to montane levels with the <i>Ranunculion fluitantis</i> and <i>Callitriche-Batrachion</i> vegetation	6510	Lowland hay meadows (<i>Altopaeurus pratensis</i> , <i>Songetisorba officinalis</i>)
3270	Rivers with muddy banks with <i>Cheopodion rubri</i> <i>p.p.</i> and <i>Bidenton p.p.</i> vegetation	7140	Transition mires and quaking bogs
3280	Constantly flowing Mediterranean rivers with <i>Paspalo-Agrostidion</i> species and hanging curtains of <i>Salix</i> and <i>Populus alba</i>	8130	Western Mediterranean and itermophilous scree
4010*	Northern Atlantic wet heaths with <i>Erica tetralix</i>	8220	Siliceous rocky slopes with chasomphytic vegetation
4020*	Temperate Atlantic wet heaths with <i>Erica ciliaris</i> and <i>Erica tetralix</i>	8230	Siliceous rock with pioneer vegetation of the <i>Sedo-Scleranthion</i> or of the <i>Sedo albi-Ieronicion dillenii</i>
4030	European dry heaths	8310	Caves not open to the public
4060	Alpine and boreal heaths	91E0*	Alluvial forests with <i>Alnus glutinosa</i> and <i>Fraxinus excelsior</i> (<i>Alno-Pandion</i> , <i>Alnion incanae</i> , <i>Salicion albae</i>)
4090	Endemic oro-Mediterranean heaths with gorse	9230	Calicio-Portuguese oak woods with <i>Quercus robur</i> and <i>Quercus pyrenaica</i>
5120	Mountain <i>Cytisus purgans</i> formations	9260	<i>Casanea sativa</i> woods
5230*	Arborecent matorral with <i>Laurus nobilis</i>	92A0	<i>Salix alba</i> and <i>Populus alba</i> galleries
5330	Thermo-Mediterranean and pre-desert scrub	9330	<i>Quercus sibirica</i> forests
6160	Oro-Iberian <i>Festuca indigesta</i> grasslands	9340	<i>Quercus flex</i> and <i>Quercus rotundifolia</i> forests
6220*	Pseudo-steppe with grasses and annuals of the <i>Thero-Brachypodetea</i>	9380	Mediterranean <i>Taxus baccata</i> woods
6230*	Species-rich <i>Nardus</i> grasslands, on siliceous substrates in mountain areas (and submountain areas, in Continental Europe)		

extremely rare groves of *Taxus baccata* (habitat 9580*). This improvement called attention to other tree vegetable communities such as the groves of holm-oaks, alder-groves and willows groves (habitats 9340, 91E0* and 92A0), certain communities typical to the upper plateau (habitats 3130, 4060 and 7140) and some communities associated to traditional practises, specifically the swamps and the *Nardus* grasslands (habitats 6230* and 6510).

The following step was the diffusion of this biological heritage. Therefore the edition of the *Guide to the Habitats of the Natural Park of Serra da Estrela* was made (Meireles *et al.*, 2006b), made in cooperation with the University of Évora. Besides that, a video for diffusion on the main habitats found in this protected area was also edited (Meireles & Neiva, 2006). These two works were destined to the general public, and intended to be an important tool to give basic knowledge of the habitats of communitarian interest found in the PNSE. These publications were possible thanks to a Life-Nature Project, which occurred between 2202 and 2006, which was called *Serra da Estrela: Gestão e Conservação de Habitats Prioritários* (management and preservation of priority Habitats). This project, which had as main goal to contribute to the recovery of priority habitats found in Serra da Estrela, had as partners the PNSE and the University of Évora, among others (Meireles *et al.*, 2004; Pinto-Gomes *et al.*, 2005, 2006).

Finally, through all these works and from the personal knowledge of the authors it was possible to define a set of specific management measures for the main habitats of this area, with the goal to preserve, recover and enhance the areas of occurrence, as well as its state of preservation.

V. PROPOSALS OF MANAGEMENT FOR HABITATS OF THE NATURAL PARK OF SERRA DA ESTRELA

Based in the works previously referred, it was possible to define measures of management for the main communitarian interest habitats found in the PNSE. In this process all the experience and results attained through the above-mentioned project Life-Nature was also fundamental. Thus, so as to propose future management measures, some geographical and inherent parameters of each one of the habitats were previously defined (distribution in Portugal, PNSE distribution, area of occurrence in the PNSE, preservation state in the PNSE and improvement). The *distribution in Portugal* was done based in the habitats files developed by the Phytosociology Lusitanian Association, in the scope of the Natura 2000 Sectorial Plan (ALFA, 2004). The *PNSE distribution* and the *area of occurrence* were made based in the above-mentioned habitat mapping, whereas the *preservation state* was made from the personal knowledge of the authors. The *area of occurrence* concerns the area where one can find the habitat and does not necessarily correspond to its area of effective occupation. Finally, the value which refers to *improvement* corresponds to the result attained in the phase of the improvement of the plant community heritage and which was defined in four categories (Exceptional, High, Average and Low) according to the preservation interest. Every time the improvement wasn't made for the habitat in question, the situation was referred to as « unknown ». The results are schematised in table II and were made for all the referenced habitats.

Next, there was the selection of the main habitats to be the target for a more profound analysis in terms of management. The selection was based in the general observation of the parameters found in table II, with particular attention to the national and local distribution, as well as the result of the improvement. Thus, 14 habitats were selected: 3130, 4010, 4060, 5230*, 6160, 6230*, 6510, 7140, 91E0*, 92A0, 9230, 9330, 9340 and 9580*.

For each of these habitats some parameters of ecological tendency and of management in reference to the PNSE (general tendency in the PNSE, main threats in the PNSE, future perspectives and management goals) were then defined. These parameters are in table III and were filled out according to the personal knowledge of the authors, having also been taken into consideration the information of the above-mentioned habitat files. The *general tendency* is in reference to the last years and the *future perspectives* are in reference to a set with no management.

Finally, the main guidance lines put forward for each of the habitats were defined, defining some sites of priority action, as well as some needed studies for their knowledge and monitoring (Table IV).

VI. CONCLUSION

The Mediterranean basin is undoubtedly an exceptional place in terms of biological variety. Just as to better explain we will stand out the fact that 24 000 species of flora, in an area

Table II - Geographical parameters, preservation state and improvement of the main habitats found in the PNSE.
 Tableau II - Paramètres géographiques, état de conservation et amélioration des principaux habitats du PNSE.

Habitat	Distribution in Portugal	Distribution in the PNSE	Area of occurrence in the PNSE	State of preservation in the PNSE	Improvement
3130	Occasional and scarce in regions but some communities only exist in the upper part of the Serra da Estrela)	Occasional. Very well represented mainly in the Upper Plateau	< 100 ha	In good state in the upper plateau. Occasionally affected by local tourism.	High
3150	Frequent	Frequent	238 km	In good state	Low
3260	Frequent	Frequent	258 km	Generally in good state	High
3270	Frequent	Unusual	16,5 km	In good state	Unknown
3280	Frequent	Unusual	16 Km	In good state	Unknown
3290	Rare (only in the tallest and rainiest mountains)	Very rare	unknown	In good state	Unknown
4010			< 10 ha	In terrible state.	High
4020	Unusual (mostly in the Centre and North)	Very rare	unknown	Present in small areas generally very altered	
4030	Very frequent	Very rare		In bad state.	High
4060	Very rare (only in the upper part of the Serra da Estrela)	Very frequent Rare (although locally very frequent over 1600 m)	27 876 ha 1 362 ha	Present in small areas generally very altered	Low
4090	Unusual (occasional in the North and Centre)	Unusual	3 829 ha	In good state	High
5120	Very rare (only in the upper part of the Serra da Estrela and Serra de Açor)	Very Rare	4 600 ha	Generally in good state	Average
5230*	Very unusual	Very Rare	33 km	Generally in good state	Average
5330	Frequent	Unusual	136 ha	Generally in terrible state of preservation with very isolated and fragmented populations.	Exceptional
6160	Rare (only in some mountainous and colder territories)	Unusual (although locally very frequent in the upper part of the Serra)	7842 ha	Generally in bad state of preservation.	Unknown
6220	Frequent (although some communities are unusual)	Unusual (although locally very frequent in the upper part of the Serra)	34 312 ha	In some spots in good state of preservation. In the areas of better access and most commonly used for touristic or recreational purposes in bad state.	Average
6230*	Unusual (only in some mountains of colder climate)	Very frequent (although some communities are rarer) Unusual (although locally frequent over 1000 metres)	6 6915 ha	In some spots in bad state	High
6410	Frequent (all over the rainier Centre and North)	Frequent	unknown	Generally in reasonable state of preservation, although locally fairly affected by natural and man action causes.	High
6430	Frequent	Frequent	unknown	In good state	Unknown
6510	Frequent in the Centre and North	Frequent	2 533 ha	In some spots in good state of preservation although in general it doesn't have the desired conditions.	High
7140	Very rare (occasional only in some mountains of the Centre and North)	Rare (although locally frequent, mainly in the Upper Plateau)	1,5 ha	(generally in a reasonable state of preservation but affected in some spots by the touristic usage of the area.	High
130	Rare	Unusual	1 593 ha	In good state	Average

of about 2,3 millions of square kilometres are referenced for this basin, that it 10% of known plants, in a small part of the world (Thompson, 2005). For the maintenance and improvement of this biodiversity it is essential to develop management measures and the preservation of its habitats, for which the Habitats Directive and its implement has greatly contributed.

This is why the PNSE strategy follows the Directive directives, developing studies in the location and mapping of the communitarian interest habitats found in the Park, as well as management actions and the recovery of the main habitats found in this Protected Area, having in mind the goal of preserving and improving biodiversity, duly associated with the economical and ecological sustenance.

Acknowledgements - We thank teacher Sandra Raposo for the translation.

REFERENCES

- ALFA, 2004. - *Tipos de Habitat Naturais e Semi-Naturais do Anexo I da Directiva 92/43/CEE (Portugal continental): Fichas de Caracterização Ecológica e de Gestão para o Plano Sectorial da Rede Natura 2000*. Relatório. Lisboa.
- Braum-Blanquet J., A.R. Pinto da Silva & A. Rozeira, 1956. - Résultats de deux excursions géobotaniques à travers le Portugal septentrional et moyen. II - Chênaies à feuilles caduques (*Quercion occidentalis*) et chênaies à feuilles persistents (*Quercion fagineae*) au Portugal. *Agron. Lusit.*, 18, 167-235.
- Braum-Blanquet J., A.R. Pinto da Silva & A. Rozeira, 1964. - Résultats de deux excursions géobotaniques à travers le Portugal septentrional et moyen. III - Landes à Cistes et Ericacées (*Cisto-Lavanduletea* et *Calluno-Ulceetea*). *Agron. Lusit.*, 23, 1229-313.
- Braum-Blanquet J., A.R. Pinto da Silva, A. Rozeira & F. Fontes, 1952. - Résultats de deux excursions géobotaniques à travers le Portugal septentrional et moyen. I - Une excursion dans la «Serra da Estrela». *Agron. Lusit.*, 14, 303-323.
- Costa J.C., C. Aguiar, J. Capelo, M. Lousã & C. Neto, 1998. - Biogeografia de Portugal Continental. *Quercetea*, 0, 5-56.
- Duvigneaud J., 1962. - Le Portugal central et septentrional. Sa position phytogéographique, sa végétation. In: *Itinéraires botaniques en Espagne et au Portugal*. L. Delvosalle & J. Duvigneaud (eds), *Les Naturalistes Belges*, 43 (2/6), 83-109.
- Esteves A., C. Meireles, J. Coimbra & C. Aguiar, 2006. - *Cartografia dos Habitats de Interesse Comunitário Presentes no Parque Natural da Serra da Estrela. Cartografia em Formato Digital*. Parque Natural da Serra da Estrela, Manteigas.
- Henriques J.A., 1883. - *Expedição científica à Serra da Estrela em 1881*. Seção de Botânica, Soc. Geogr., Lisboa.
- Jansen J., 1997. - *A survey of habitats and species occurring in the Parque Natural da Serra da Estrela*. Final report for Natura 2000 Project. Museu e Jardim Botânico, Universidade de Lisboa, 137 p.
- Jansen J., 2002. - *Guia geobotânico da Serra da Estrela*. Parque Natural da Serra da Estrela, Manteigas, 276 p.
- Maeza G. & J.A. Cadizanos, 2000. - *Valoración de la Vegetación*. In: *Metodología y práctica de la biogeografía*. Ediciones del Serbal, 201-272.
- Maliato-Belz J., 1955. - *As pastagens de servum (Nardus stricta L.) da Serra da Estrela: fitossociologia e mel-*

Table III.- Ecological tendency and management parameters, for the PNSE, of the main habitats found in this protected area.
Tableau III.- Tendence écologique et paramètres de gestion des principaux habitats de l'aire protégée du PNSE.

Habitat	General tendency	Main threats	Perspectives for the future	Management goal(s)
3130	Decrease	- Water dystroficaton - Direct destruction - Disorderly tourism - Drainage - Global warming	Decrease of the area of occurrence	- Maintenance of the area of occurrence - Encouragement of the <i>Spartanium angustifolium</i> communities
4010	Decrease	- Fires - Direct destruction for the alteration of soil usage - Alteration of the hydric regimen	Decrease of the area of occurrence and maintenance of the preservation degree	- Encouragement of the area of occurrence - Improvement of the preservation state
4060	Decrease	- Direct destruction mostly by the creation of touristic support substructures (mostly associated to sky)	Maintenance/decrease of the area of occurrence and occasional deterioration of the state of preservation	- Maintenance of the current area of occurrence and of the state of preservation
5230	Decrease	- Fires - Low cut - Invasion by exotic species	Decrease	- Increase of the current area of occurrence for twice the present
5160	Decrease	- Physical destruction for the construction of substructures - Decrease of grazing pressure	Decrease	- Preservation of the current area of occurrence - Improvement of the state of preservation
230	Decrease	- Desertion of agro-sylvo-pastoral activities - Successional progression - Soil mobilization (in areas of less altitude)	Decrease	- Increase of the area occupied by the habitat - improvement of the preservation state
510	Decrease	- Desertion of traditional agro-sylvo-pastoral activities and practices - Alteration in the usage of the soil	Decrease	- Maintenance of the area occupied by the habitat - Improvement of the general state of preservation
140	Decrease	- Physical destruction - Water dystroficaton	Maintenance	- Maintenance of the area occupied by the habitat - Improvement of the general state of preservation
1E0	Decrease	- Low cut - Alteration of the hydric regimen	Maintenance	- Encouragement of the area of occurrence - Improvement of the general state of preservation
2A0	Decrease	- Low cut - Fires	Maintenance	- Encouragement of the area of occurrence - Improvement of the general state of preservation
30	Decrease	- Fires	Decrease	- Encouragement of the area of occurrence - Improvement of the general state of preservation - Fire protection
30	Maintenance	- Low cut - Fires	Maintenance /Increase	- Encouragement of the area of occurrence

9140	Decrease	- Low cut - Fires	Maintenance	- Improvement of the general state of preservation - Fire protection
9580	Decrease	- Fires	Decrease	- Encouragement of the area of occurrence - Improvement of the general state of preservation - Fire protection

horamento. *Melhoramento*, VIII, 59 p.

Meireles C. & R. Neiva, 2006.- *Habitats do Parque Natural da Serra da Estrela*. Vídeo divulgativo, Parque Natural da Serra da Estrela, Manteigas.

Meireles C., R. Neiva, A. Correia & C. Pinto-Gomes, 2006b.- *Guia de habitats do Parque Natural da Serra da Estrela*. Parque Natural da Serra da Estrela, Manteigas, 67 p.

Meireles C., R. Paiva-Ferreira & C. Pinto-Gomes, 2004.- *Serra da Estrela; gestão e conservação de habitats prioritários (Life02 nat/p/8478)*. 1º relatório de progresso, Universidade de Évora, 68 p.

Meireles C., C. Pinto-Gomes, R. Paiva-Ferreira & E. Cano, 2006a.- *Valorização do território do Parque Natural da Serra da Estrela, desde o ponto de vista da flora e da vegetação*. Apresentação oral e em poster. IV Congresso Espanhol de Biogeografia, Ávila.

Passos I., 2005.- *Habitats naturais e semi-naturais do Vale Glaciar do Rio Zêzere (Serra da Estrela)*; cartografia pré e pós-incêndio de 2005. Trabalho de fim de curso em Biologia, Universidade de Évora.

Pinto da Silva A.R. & A.N. Teles, 1986.- *A flora e a vegetação da Serra da Estrela. Coleção Parques Naturais*, 7 (2ª Edição), 52 p.

Pinto da Silva A.R., A.N. Teles & M.H.R. Lopes, 1976.- *Teucrium salvivstrum* Schreb. *Agron. Lusit.*, **37**, 184.

Pinto-Gomes C. & R. Paiva-Ferreira, 2005.- *Serra da Estrela; gestão e conservação de habitats prioritários (Life02 nat/p/8478)*. 2º relatório de progresso. Universidade de Évora, 44 p.

Pinto-Gomes C. & R. Paiva-Ferreira, 2006.- *Serra da Estrela; gestão e conservação de habitats prioritários (Life02 nat/p/8478)*. Relatório final, Universidade de Évora, 156 p.

Rivas-Martínez S., 1974.- *Datos sobre la flora y la vegetación de la Serra da Estrela (Portugal)*. *Anal. R. Acad. Farm.*, **40** (1), 65-74.

Rivas-Martínez S., 1987.- *Mapa de series de vegetación de España*. ICONA, Serie técnica, Madrid, 268 p. + 30 mapas.

Rivas-Martínez S., T.E. Diaz, F. Fernández-González, J. Izco, X. Loidi, M. Lousã & A. Penas, 2002.- *Vascular plant communities of Spain and Portugal*. addenda to the syntaxonomical checklist of 2001. *Itiner Geobot.*, **15** (1-2), 1-922.

Sampaio G., 1910.- *Flora das lagoas*. In: *Notas sobre a fauna das águas das lagoas da Serra da Estrela*. A. Nobre (ed.), *Bol. Dir. Ger. Agric.*, **9** (7), 1-7.

Thompson J., 2005.- *Plant evolution in the Mediterranean*. Oxford University Press, 293 p.

ole IV.- Management measures for the main habitats found in the PNSE.
bleau IV.- Mesures de gestion sur les principaux habitats du PNSE.

Proposed management measures	Places of intervention	Parameters to monitor / study
<ul style="list-style-type: none"> - Interdiction to swimming in the ponds of the upper plateau - Spreading of the <i>Spartanium angustifolium</i> to ponds from where it has disappeared. 	Upper Plateau	<ul style="list-style-type: none"> - Water quality - Communities of <i>Spartanium angustifolium</i> - Population of <i>Menyanthes trifoliata</i>
<ul style="list-style-type: none"> - Insuring the maintenance of the soil usage - Implementing the selective cut of vegetation to stop the natural evolution into upper steps - Rectify the water drainage when necessary - Protecting the areas against fires (mainly at higher altitude) 	Upper Plateau and "Unhais" Valley	<ul style="list-style-type: none"> - Thorough inventory of the cores and of their state of preservation
<ul style="list-style-type: none"> - Insuring the non-expansion of substructures for its area of occurrence - Stopping the traffic in its areas of occurrence 	Upper Plateau	<ul style="list-style-type: none"> - Extending the area occupied by the habitat - State of preservation - Population of <i>Lycopodium clavatum</i>
<ul style="list-style-type: none"> - Protection of the areas against fires - Recovery of the areas affected by fires - Extirmination of the exotic species of its areas of occurrence (mostly of <i>Pinus pinaster</i>, <i>Eucalyptus</i> sp.) 	"Casal do Rei" and "Cabeça"	<ul style="list-style-type: none"> - Thorough inventory of the cores and of their state of preservation - Recovery of the post-fire areas
<ul style="list-style-type: none"> - Insuring physical maintenance - Encouraging pastoral activity 	Upper Plateau	<ul style="list-style-type: none"> - Capability of recovery of the areas after physical destruction
<ul style="list-style-type: none"> - Selective cut - Installing small walls - Encouraging pastoral activity - Controlling the alteration in soil usage ("Videmonte") 	Upper Plateau and "Videmonte" Plateau	<ul style="list-style-type: none"> - Area of occupation (with detailed mapping)
<ul style="list-style-type: none"> - Encouraging traditional agro-sylvo-pastoral practices 	Mainly Glacier of the Valley of the "Zézere" River, "Mondego" and "Videmonte" Plateau	<ul style="list-style-type: none"> - Occupation area
<ul style="list-style-type: none"> - Restoring favourable ecological conditions - Avoiding water drainage - Avoiding the use of fire in these areas 	Upper Plateau	<ul style="list-style-type: none"> - Water quality
<ul style="list-style-type: none"> - Avoiding the low cut - Encouraging planting 	Along "Zézere" Valley	<ul style="list-style-type: none"> - Occupation area
<ul style="list-style-type: none"> - Avoiding low cut - Planting endemic willows in places where they disappear from 	All over the Park	<ul style="list-style-type: none"> - Occupation area
<ul style="list-style-type: none"> - Avoiding low cut - Fighting fires 	In the few existent spots in the "PNSE"	<ul style="list-style-type: none"> - Occupation area
<ul style="list-style-type: none"> - Avoiding low cut - Fighting fires 	In the few existent spots in the "PNSE"	<ul style="list-style-type: none"> - Occupation area
<ul style="list-style-type: none"> - Avoiding low cut - Fighting fires 	In the few existent spots in the "PNSE"	<ul style="list-style-type: none"> - Occupation area
<ul style="list-style-type: none"> - Avoiding low cut - Fighting fires 	"Zézere" Valley "Caniceal" Stream "Bejames" Stream	<ul style="list-style-type: none"> - Occupation area - Recovery after fire - Growth rate - Genetic diversity of <i>Tarax baccata</i>

Contribution to the knowledge of plant diversity and conservation of natural areas in la Manchuela Conquense (Spain)

by Maria Pilar Rodriguez-Rojo and Rosa Pérez-Badia

Area of Botany, Environmental Sciences Institute, Faculty of Environmental Sciences, University of Castile-La Mancha, E-45071 Toledo, mpilar.rodriguez@uclm.es

Abstract. - This paper reports the results of a floristical and phytosociological study on La Manchuela Conquense, a region in Castile-La Mancha (Spain) characterized by a multiple complex of biotopes which sustain a great floristical and community diversity. More than 60% of its species are Mediterranean, in which an important percentage are endemics. Approximately 11% of the total of species are Iberian-Magreb, Iberian or lower rank endemics. Levantine-Iberians contribute in majority, to total endemicity, but there is a group of rare species living on siliceous soils which contribute to the floristic diversity of this territory. A short description of community diversity is also included attending to vegetation series. Based on the distribution of endangered and singular species and habitat in the territory, we may have brought out some areas of natural value to be considered for management and conservation planning.

Key words : Cuenca - flora - habitat - phytosociology - endangered species.

Résumé. - Ce papier rapporte les résultats d'une étude floristique et phytosociologique menée dans la Manchuela Conquense, une région de Castille-La Mancha (Espagne) caractérisée par un complexe de biotopes avec une grande diversité de communautés végétales. Plus de 60% des espèces sont méditerranéennes, avec un fort pourcentage d'endémiques. Près de 11% du total des espèces sont endémiques de la péninsule ibérique et du Maghreb ou d'un niveau inférieur. Les espèces levantines-ibériques contribuent en grande partie à l'endémicité totale, mais on relève un rare groupe d'espèces dans les terrains siliceux qui contribue à la diversité floristique de ce territoire. Une brève description de la diversité des communautés est aussi abordée sur la base des séries de végétation. Basée sur la distribution d'espèces particulières en danger et l'habitat dans le territoire, nous croyons distingué montré plusieurs aires naturelles devant être intégrées dans la planification de sa gestion et sa conservation.

Mots clés : Cuenca - flora - habitat - phytosociologie - espèces menacées.