



## Reconstructing the architectural evolution of volcanic islands from combined K/Ar, morphologic, tectonic, and magnetic data: The Faial Island example (Azores)

A. Hildenbrand<sup>a,b,\*</sup>, F.O. Marques<sup>c</sup>, A.C.G. Costa<sup>a,c</sup>, A.L.R. Sibrant<sup>a,c</sup>, P.F. Silva<sup>d</sup>, B. Henry<sup>e</sup>, J.M. Miranda<sup>c</sup>, P. Madureira<sup>f,g</sup>

<sup>a</sup> Univ Paris-Sud, Laboratoire IDES, UMR8148, Orsay, F-91405, France

<sup>b</sup> CNRS, Orsay, F-91405, France

<sup>c</sup> Universidade de Lisboa, IDL, Lisboa, Portugal

<sup>d</sup> ISEL/DEC and IDL/CGUL, Lisboa, Portugal

<sup>e</sup> Paleomagnetism, IPGP and CNRS, 4 Av. de Neptune, 94107 Saint-Maur cedex, France

<sup>f</sup> Centro de Geofísica de Évora and Dep. de Geociências da Univ. de Évora, R. Romão Ramalho, 59, 7000-671 Évora, Portugal

<sup>g</sup> Estrutura de Missão para os Assuntos do Mar, R. Costa Pinto, 165, 2770-047, Paço D'Arcos, Portugal

### ARTICLE INFO

#### Article history:

Received 10 January 2012

Accepted 11 June 2012

Available online 21 June 2012

#### Keywords:

Faial

Volcanic construction

Mass-wasting

K/Ar

Tectonics

Magnetism

Azores triple junction

### ABSTRACT

The morpho-structural evolution of oceanic islands results from competition between volcano growth and partial destruction by mass-wasting processes. We present here a multi-disciplinary study of the successive stages of development of Faial (Azores) during the last 1 Myr. Using high-resolution digital elevation model (DEM), and new K/Ar, tectonic, and magnetic data, we reconstruct the rapidly evolving topography at successive stages, in response to complex interactions between volcanic construction and mass wasting, including the development of a graben. We show that: (1) sub-aerial evolution of the island first involved the rapid growth of a large elongated volcano at ca. 0.85 Ma, followed by its partial destruction over half a million years; (2) beginning about 360 ka a new small edifice grew on the NE of the island, and was subsequently cut by normal faults responsible for initiation of the graben; (3) after an apparent pause of ca. 250 kyr, the large Central Volcano (CV) developed on the western side of the island at ca 120 ka, accumulating a thick pile of lava flows in less than 20 kyr, which were partly channelized within the graben; (4) the period between 120 ka and 40 ka is marked by widespread deformation at the island scale, including westward propagation of faulting and associated erosion of the graben walls, which produced sedimentary deposits; subsequent growth of the CV at 40 ka was then constrained within the graben, with lava flowing onto the sediments up to the eastern shore; (5) the island evolution during the Holocene involves basaltic volcanic activity along the main southern faults and pyroclastic eruptions associated with the formation of a caldera volcano-tectonic depression. We conclude that the whole evolution of Faial Island has been characterized by successive short volcanic pulses probably controlled by brief episodes of regional deformation. Each pulse has been separated by considerable periods of volcanic inactivity during which the Faial graben gradually developed. We propose that the volume loss associated with sudden magma extraction from a shallow reservoir in different episodes triggered incremental downward graben movement, as observed historically, when immediate vertical collapse of up to 2 m was observed along the western segments of the graben at the end of the Capelinhos eruptive crises (1957–58).

© 2012 Elsevier B.V. All rights reserved.

### 1. Introduction

The geological evolution of oceanic islands is generally marked by rapid volcanic growth alternating with destruction by a variety of mass-wasting processes including giant sector collapses, vertical caldera subsidence, fault generation/propagation, shallow landslides and coastal erosion (e.g., Moore et al., 1989; Mitchell, 1998; Hildenbrand

et al., 2004, 2006, 2008a, 2008b; Quartau et al., 2010; Boulesteix et al., 2012). The study of such destruction events is of particular societal relevance, since they can cause considerable damage. The Azores volcanic islands in the Atlantic are particularly sensitive to mass-wasting processes. The region is characterized by intense deformation responsible for high magnitude earthquakes (e.g., Borges et al., 2007). Like the neighbouring islands of Pico, S. Jorge, Terceira and Graciosa in the central Azores, Faial is still active and is characterized by the localization of volcanism along N110°-trending eruptive fissures and vents. Faial is additionally marked by a large graben-like structure elongated along the same azimuth (N110°). The island thus offers a great opportunity

\* Corresponding author at: Univ Paris-Sud, Laboratoire IDES, UMR8148, 91405 Orsay, F-91405, France. Tel.: +33 1 69 15 67 42; fax: +33 1 69 15 48 91.

E-mail address: [anthony.hildenbrand@u-psud.fr](mailto:anthony.hildenbrand@u-psud.fr) (A. Hildenbrand).